arXiv:1212.5131v1 [hep-th] 20 Dec 2012

Noncommutative field theories on $\mathbb{R}^3_\lambda$: Toward UV/IR mixing freedom

Patrizia Vitale [a,b,c] and Jean-Christophe Wallet[c]

[a]*Dipartimento di Scienze Fisiche Università di Napoli Federico II*
[b]*INFN, Sezione di Napoli, Via Cintia 80126 Napoli, Italy*
[c]*Laboratoire de Physique Théorique, Bât. 210*
CNRS and Université Paris-Sud 11, 91405 Orsay Cedex, France

e-mail: `patrizia.vitale@na.infn.it`, `jean-christophe.wallet@th.u-psud.fr`

Abstract

We consider the noncommutative space $\mathbb{R}^3_\lambda$, a deformation of the algebra of functions on $\mathbb{R}^3$ which yields a "foliation" of $\mathbb{R}^3$ into fuzzy spheres. We first construct a natural matrix base adapted to $\mathbb{R}^3_\lambda$. We then apply this general framework to the one-loop study of a two-parameter family of real-valued scalar noncommutative field theories with quartic polynomial interaction, which becomes a non-local matrix model when expressed in the above matrix base. The kinetic operator involves a part related to the square of the angular momentum supplemented by a term representing radial dynamics. We then compute the planar and non-planar 1-loop contributions to the 2-point correlation function. When only the angular momentum part of the kinetic operator is involved, the planar 1-loop contribution is logarithmically divergent while the non-planar contribution has a soft behavior in the external momenta, which signals very likely the absence of UV/IR mixing. We also consider the case of a kinetic operator with only the "radial part". We find that the resulting theory is finite to all orders in perturbation expansion.

1 Introduction.

Noncommutative Geometry (NCG) [1] provides a generalization of topology, differential geometry and index theory. The starting idea is to set-up a duality between spaces and associative algebras in a way to obtain an algebraic description of the structural properties of the space, in particular the topological, metric, differential properties [2]. Besides, many of the building blocks of fundamental physics fit well with concepts of NCG which may lead to a more accurate understanding of spacetime at short distance and/or possibly of what could be a quantum theory of gravity. For instance, NCG offers a possible way to treat the physical obstructions to the existence of a continuous space-time and commuting coordinates at the Planck scale [3]. Once the noncommutative nature of space-time postulated, it is natural to consider field theories on noncommutative manifolds, the so called Noncommutative Field Theories (NCFT).

The first prototypes of NCFT appeared in 1986 within String field theory [4]. Field theories defined on the fuzzy sphere, a simple finite dimensional noncommutative geometry [5, 6, 7], were introduced at the beginning of the 90's in [8] and actively studied since then. See for example [9] for a review. In1998 NCFT on the Moyal space, the simplest noncommutative geometry modeled on the phase-space of quantum mechanics, was shown to occur in effective regimes of String theory [10]. This observation triggered a huge activity. Noncommutative field theory of Moyal type was also shown to describe quite accurately quantum Hall physics [11]. For a review on NCFT on Moyal spaces see [12, 13] and references therein.

The renormalization study of NCQFT is in general difficult, a part from the case of finite noncommutative geometries, and is often complicated by the Ultraviolet/Infrared (UV/IR) mixing. This phenomenon occurs for instance within the simplest noncommutative real-valued φ^4 model on the 4-dimensional Moyal space, as pointed out and analyzed in [14]. The phenomenon persists in Moyal-noncommutative gauge models and represents one of the main open problems of Moyal-based field theory. In [15] noncommutative differential structures relevant to Moyal-noncommutative gauge theories were studied, precisely to tackle such a problem. A first solution for scalar field theory was proposed in 2003 [16]. It amounts to modify the initial action with a harmonic oscillator term leading to a fully renormalisable NCFT. This is the so called Grosse-Wulkenhaar model. Various of its properties have been explored, among which classical and/or geometrical ones [17], the 2-d fermionic extension [18] as well as the generalization to gauge theory (matrix) models [19], [20], [21], [22]. The Grosse-Wulkenhaar model has interesting properties such as the vanishing of the β-function to all orders [23] when the action is self-dual under the so-called Langmann-Szabo duality [24]. Its 4-d version is very likely to be non-perturbatively solvable, as shown in [25]. Besides, this model together with its gauge theory counterpart seems to be related to an interesting noncommutative structure, namely a finite volume spectral triple [26] whose relation to the Moyal (metric) geometries has been analyzed in [27, 28].

This paper deals with a different kind of noncommutativity, said of "Lie algebra" type, because the $\star$-commutator of coordinate functions is not constant and reproduces the Lie bracket of classical Lie algebras. We shall follow [29] where many $\star$-products were proposed, reproducing at the level of coordinate functions all three-dimensional Lie algebras, and in particular [30], where the specific case of a $\mathfrak{su}(2)$ based start product giving raise to the noncommutative space $\mathbb{R}^3_\lambda$ was first introduced. The purpose of this paper is twofold. The first goal is to set-up a general framework that can be used to study the quantum (i.e renormalisability) behaviour of matter NCFT as well as gauge NCFT [31] defined on $\mathbb{R}^3_\lambda$. The second goal is to apply this framework to a class of natural scalar NCFT on $\mathbb{R}^3_\lambda$ in order to capture salient information related to its one-loop behaviour. There are important differences between $\mathbb{R}^3_\lambda$ and the popular Moyal space. First of all, the $\star$-commutator between the coordinates of $\mathbb{R}^3_\lambda$ is no longer constant and the relevant algebra of functions coding the $\mathbb{R}^3_\lambda$ NCG is equipped with an associative but not *translation-invariant* product. Moreover, the popular tracial property of the Moyal algebra [32] does not hold true, which complicates the treatment of the kinetic part of the action. This difficulty can be handled by using a suitable matrix base. This is one of the results of the paper. We construct a natural matrix base adapted to $\mathbb{R}^3_\lambda$ which can be obtained as a reduction of the matrix base of the Moyal space $\mathbb{R}^4_\theta$ [32][*]. We then consider a family of real-valued scalar actions on $\mathbb{R}^3_\lambda$ with quartic polynomial interactions. The family of kinetic operators, indexed by two real parameters, involves a natural Laplacian-type operator which contains the square of the angular momentum and a radial part related (but not equal) to the Casimir operator of $\mathfrak{su}(2)$, which generates some radial dynamics. When re-expressed in the natural matrix base, the action is the one for a non-local matrix model with *diagonal* interaction term[†] and *non-diagonal* kinetic operator being of Jacobi type. The action can be split as an infinite sum of scalar actions defined on the successive fuzzy spheres that "foliate" the noncommutative space $\mathbb{R}^3_\lambda$. The radial term in the Laplacian encodes the radial dynamics between spheres. Another matrix base, largely used in the literature on the fuzzy sphere, is built from the symbols of the fuzzy spherical harmonic operators and related to the natural base of $\mathbb{R}^3_\lambda$. This leaves *diagonal* the kinetic operator with however a complicated interaction term[‡]. Upon diagonalizing the kinetic term, the computation of the propagator in the natural base can then be performed and we end up with a tractable expression together with a purely diagonal interaction term. We then compute the planar and non-planar 1-loop contribution to the 2-point correlation function. When only the angular momentum part of the kinetic operator is involved, the resulting scalar action leads to a logarithmically divergent planar 1-loop contribution. Using asymptotic expressions for the Wigner $6j$-coefficients entering the non-planar contribution, we find that the latter has a soft behaviour in the external momenta (indices). This signals very likely the absence of the UV/IR mixing phenomenon at the perturbative level. We also consider the other limit situation with a kinetic operator without angular momentum part and only the "radial part". This leads to a big simplification for the action and for the general power counting of the ribbon diagrams of arbitrary orders. We find that the resulting theory is finite to all orders in perturbation.

The paper is organized as follows. In section 2 we summarize the general properties of the noncommutative $\mathbb{R}^3_\lambda$ that will be used in this paper together with some features related to the Voros product. In section 3 we construct a natural matrix base adapted to $\mathbb{R}^3_\lambda$. In section 4 we construct a family of real-valued scalar actions on $\mathbb{R}^3_\lambda$ with quartic polynomial interactions. The relationship with the base built from the fuzzy spherical harmonics is also introduced. The subsection 4.3 involves the computation of the propagator expressed in the natural matrix base for $\mathbb{R}^3_\lambda$, which is rather easily carried out, once a suitable combination of the change of base of subsection 4.2 with properties of fuzzy spherical harmonics is done. Nevertheless, we find interesting to provide in the appendix the general computation of the propagator that takes advantage of the Jacobi nature of the kinetic operator. This is based on the determination of a suitable family of orthogonal polynomials that gives rise to diagonalization. We find in the present case that the relevant orthogonal polynomials are the dual Hahn polynomials, the counterpart of the Meixner polynomials underlying the computation of the Grosse-Wulkenhaar propagator. This, as a by-product, provides explicit relations between fuzzy spherical harmonics, Wigner $3j$-symbols and dual Hahn polynomials. In section 5 we compute and discuss the planar and non-planar 1-loop contribution to the 2-point correlation function, respectively in the subsections 5.1 and 5.2. In the subsection 5.3 we consider the other limit case of a kinetic operator with no angular momentum term and only the radial term, for which we find finitude to all orders in perturbation. We finally summarize the results and conclude.

---

[*] As we shall see in detail the starting matrix base in $\mathbb{R}^4_\theta$ is actually a slight modification of the Moyal matrix base which was introduced in [32]. We shall use a matrix base adapted to the Voros product.

[†] In physical words, the natural base is nothing but the interaction base.

[‡] This other base is physically the so-called propagation base. See the previous footnote. We will use this terminology when appropriate in the paper.

2 The noncommutative space $\mathbb{R}^3_\lambda$

The noncommutative space $\mathbb{R}^3_\lambda$ has been first introduced in [30]. A generalization has been studied in [29]. It is a subalgebra of $\mathbb{R}^4_\theta$, the noncommutative algebra of functions on $\mathbb{R}^4 \simeq \mathbb{C}^2$ endowed with the Voros product [33]

$$\phi \star \psi\,(z_a, \bar{z}_a) = \phi(z, \bar{z}) \exp(\theta \overleftarrow{\partial}_{z_a} \overrightarrow{\partial}_{\bar{z}_a}) \psi(z, \bar{z}), \quad a = 1, 2 \tag{2.1}$$

This is an asymptotic expansion; a proper definition, based on the dequantization map associated to normal ordered quantization,will be given in sect. 3.1 where it is actually needed to introduce a matrix base. For coordinate functions we have the $\star$-commutator

$$[z_a, \bar{z}_b]_\star = \theta \delta_{ab} \tag{2.2}$$

with θ a constant, real parameter. Resorting to real coordinates $q_a = z_a + \bar{z}_a, p_a = i(z_a - \bar{z}_a)$, $a = 1, 2$ we recover the usual $\star$-commutator of (two copies of) the Moyal plane, the two products differing by symmetric terms.[§] The crucial step to obtain star products on $\mathcal{F}(\mathbb{R}^3)$, hence to deform $\mathcal{F}(\mathbb{R}^3)$ into a noncommutative algebra, is to identify $\mathbb{R}^3$ with the dual, $\mathfrak{g}^*$, of some chosen three dimensional Lie algebra $\mathfrak{g}$. We choose here to work with the $\mathfrak{su}(2)$ Lie algebra, because of the connection with other results already present in the literature (for example the fuzzy sphere) but other choices can be made. This identification induces on $\mathcal{F}(\mathbb{R}^3)$ the Kirillov Poisson bracket, which, for coordinate functions reads

$$\{x_i, x_j\} = c_{ij}^k x_k \tag{2.3}$$

with $i = 1, .., 3$ and c_{ij}^k the structure constants of $\mathfrak{su}(2)$. On the other hand, it is well known that this (Poisson) Lie algebra may be regarded as a subalgebra of the symplectic algebra $\mathfrak{sp}(4)$, which is classically realized as the Poisson algebra of quadratic functions on $\mathbb{R}^4$ ($\mathbb{C}^2$ with our choices) with canonical Poisson bracket

$$\{z^a, \bar{z}^b\} = i. \tag{2.4}$$

Indeed it is possible to find quadratic functions

$$\pi^*(x_i) = \pi^*(x_i)(z^a, \bar{z}^a) \tag{2.5}$$

which obey (2.3).We have indicated with π^* the pull-back map $\pi^* : \mathcal{F}(\mathbb{R}^3) \to \mathcal{F}(\mathbb{R}^4)$. This is nothing but the classical counterpart of the Jordan-Schwinger map realization of Lie algebra generators in terms of creation and annihilation operators [39]. Then one can show that this Poisson subalgebra is also a Voros subalgebra, that is

$$\pi^*(x_i)(z^a, \bar{z}^a) \star \pi^*(x_j)(z^a, \bar{z}^a) - \pi^*(x_j)(z^a, \bar{z}^a) \star \pi^*(x_i)(z^a, \bar{z}^a) = \lambda c_{ij}^k \pi^*(x_k)(z^a, \bar{z}^a) \tag{2.6}$$

where the noncommutative parameter λ shall be adjusted according to the physical dimension of the coordinate functions x_i. We shall indicate with $\mathbb{R}^3_\lambda$ the noncommutative algebra $(\mathcal{F}(\mathbb{R}^3), \star)$. Eq. (2.6) induces a star product on polynomial functions on $\mathbb{R}^3$ generated by the coordinate functions

---

[§] The Moyal and Voros algebras are isomorphic [34], there is however a debate on the physical meaning of the equivalence between them, which we will not address here (see for example [35, 36, 37, 38]).

x_i, which may be expressed in closed form in terms of differential operators on $\mathbb{R}^3$. Here we will consider quadratic realizations of the kind

$$\pi^*(x_\mu) = \frac{\lambda}{\theta} \bar{z}^a e_\mu^{ab} z^b, \quad \mu = 0, .., 3 \tag{2.7}$$

with λ a constant, real parameter of length dimension equal to one; $e_i = \frac{1}{2}\sigma_i$, $i = 1, .., 3$ are the $SU(2)$ generators in the defining representation and σ_i are the Pauli matrices, while $e_0 = \frac{1}{2}\mathbf{1}$. We shall omit the pull-back map from now on, unless necessary. Notice that

$$x_0 = \frac{\lambda}{2\theta} \bar{z}_a z_a \tag{2.8}$$

commutes with x_i so that we can alternatively define $\mathbb{R}^3_\lambda$ as the commutant of x_0; x_0 generates the center of the algebra. We also have

$$x_0^2 = \sum_i x_i^2. \tag{2.9}$$

It is easily verified that the induced $\star$-product reads

$$\phi \star \psi\,(x) = \exp\left[\frac{\lambda}{2}\left(\delta^{ij} x^0 + i\epsilon^{ijk} x^k\right) \frac{\partial}{\partial u^i} \frac{\partial}{\partial v^j}\right] \phi(u)\psi(v)|_{u=v=x} \tag{2.10}$$

which implies, for coordinate functions

$$x^i \star x^j \quad = \quad x^i x^j + \frac{\lambda}{2}\left(x^0 \delta ij + i\epsilon^{ij}_k x^k\right) \tag{2.11}$$

$$x^0 \star x^i \quad = \quad x^i \star x^0 = x^0 x^i + \frac{\lambda}{2} x^i \tag{2.12}$$

$$x_0^{*2} \quad = \quad x_0(x_0 + \frac{\lambda}{2}). \tag{2.13}$$

where we have used (2.9). As for the $\star$ commutator we have

$$[x_i, x_j]_\star = i\lambda\epsilon_{ij}^k x_k. \tag{2.14}$$

On introducing the parameter $\kappa = \lambda/\theta$, the commutative limit is achieved with $\lambda, \theta \to 0$. $\kappa = const.$

We have thus realized the announced isomorphism between the algebra of linear functions on $\mathbb{R}^3 \simeq \mathfrak{su}(2)^*$ endowed with the $\star$ commutator (2.14) and the $\mathfrak{su}(2)$ Lie algebra. Thus, the algebra $\mathbb{R}^3_\lambda$ can be defined as $\mathbb{R}^3_\lambda = \mathbb{C}[x_\mu]/\mathcal{I}_{\mathcal{R}_1,\mathcal{R}_2}$, i.e the quotient of the free algebra generated by the coordinate functions $(x_i)_{i=1,2,3}, x_0$, by the two-sided ideal generated by the relation $\mathcal{R}_1 : [x_i, x_j]_\star = i\lambda\epsilon_{ijk} x_k$, together with $\mathcal{R}_2 : x_0 \star x_0 - \lambda x_0 = x_i \star x_i$. Notice that, because of the presence of x_0 $\mathbb{R}^3_{\lambda\neq 0}$ *is not* isomorphic to $\mathcal{U}(\mathfrak{su}(2))$.

For a comparison with the Moyal induced product we refer to [29].

3 The matrix base

The matrix base we shall define for $\mathbb{R}^3_\lambda$ is obtained through a suitable reduction of the matrix base of the Voros algebra $\mathbb{R}^4_\theta$, which was introduced in [40]. The latter is in turn a slight modification of the well known matrix base for the Moyal algebra defined in [32] by Gracia-Bondìa and Varilly.

3.1 The matrix base for the Voros $\mathbb{R}^4_\theta$

Let us first review the matrix base adapted to the Voros algebra $\mathbb{R}^4_\theta$. Our convention, all over the paper, will be to use hatted letters to indicate operators and un-hatted ones to indicate their noncommutative symbols.

It is well known that the Voros product is introduced through a weighted quantization map which, in two dimensions, associates to functions on the complex plane the operator

$$\hat{\phi} = \hat{\mathcal{W}}_V(\phi) = \frac{1}{(2\pi)^2}\int \mathrm{d}^2 z\, \hat{\Omega}(z,\bar{z})\phi(z,\bar{z}) \tag{3.1}$$

where

$$\hat{\Omega}(z,\bar{z}) = \int \mathrm{d}^2\eta\, e^{-(\eta\bar{z}-\bar{\eta}z)} e^{\theta\eta a^\dagger} e^{-\theta\bar{\eta}a} \tag{3.2}$$

is the so called quantizer and $a, a^\dagger$ are the usual (configuration space) creation and annihilation operators, with commutation relations

$$[a, a^\dagger] = \theta. \tag{3.3}$$

A word of caution is in order, concerning the domain and the range of the weighted Weyl map in Eq. (3.1). While the standard Weyl map maps Schwarzian functions into Hilbert Schmidt operators, for the weighted Weyl map (3.1) this is not always the case. An exhaustive analysis is lacking in the literature, up to our knowledge. Explicit counterexamples are discussed in [40].

The inverse map which is the analogue of the Wigner map is represented by:

$$\phi(z,\bar{z}) = \mathcal{W}_V^{-1}(\hat{\phi}) = \langle z|\hat{\phi}|z\rangle \tag{3.4}$$

with $|z\rangle$ the *coherent states* defined by $a|z\rangle = z|z\rangle$. Notice that, differently from the Weyl-Wigner-Moyal case, the quantizer and dequantizer operators do not coincide, meaning that this quantization/dequantization procedure is not self-dual (see [32, 41] for details).

The Voros product, whose asymptotic form has been already given in (2.1), is then defined as

$$\phi \star \psi := \mathcal{W}_V^{-1}\left(\hat{\mathcal{W}}_V(\phi)\hat{\mathcal{W}}_V(\psi)\right) = \langle z|\hat{\phi}\,\hat{\psi}|z\rangle \tag{3.5}$$

It can be seen that, for analytic functions, a very convenient way to reformulate the quantization map (3.1) is to consider their analytic expansion

$$\phi(\bar{z}, z) = \sum_{pq} \tilde{\phi}_{pq}\bar{z}^p z^q\,, \quad p, q \in \mathbb{N} \tag{3.6}$$

with $\tilde{\phi}_{pq} \in \mathbb{C}$. The quantization map (3.1) will then produce the normal ordered operator

$$\hat{\phi} = \hat{\mathcal{W}}_V(\phi) = \sum_{pq} \tilde{\phi}_{pq} a^{\dagger p} a^q \tag{3.7}$$

We will therefore assume analyticity all over the paper and use (3.6), (3.7).

This construction may be easily extended to $\mathbb{R}^{2n}$. We will consider $n = 2$ from now on.

Each a_a, $a = 1, 2$, acts on $\mathcal{H}_0 \cong \ell^2(\mathbb{N})$, a copy of the Hilbert space of the one dimensional harmonic oscillator with canonical orthonormal base $(|n\rangle)_{n\in\mathbb{N}}$. We set

$$|N\rangle = |n_1, n_2\rangle := |n_1\rangle \otimes |n_2\rangle \tag{3.8}$$

the canonical orthonormal base for $\mathcal{H} = \mathcal{H}_0 \otimes \mathcal{H}_0$, also called in the physics literature the number base. The action of the $a_a, a_a^\dagger$'s on $\mathcal{H}$ is given by

$$\begin{aligned} a_1|n_1, n_2\rangle &= \sqrt{\theta}\sqrt{n_1}|n_1 - 1, n_2\rangle, \ a_1^\dagger|n\rangle = \sqrt{\theta}\sqrt{n_1+1}|n_1+1, n_2\rangle, \\ a_2|n_1, n_2\rangle &= \sqrt{\theta}\sqrt{n_2}|n_1, n_2 - 1\rangle, \ a_2^\dagger|n\rangle = \sqrt{\theta}\sqrt{n_2+1}|n_1, n_2+1\rangle. \end{aligned} \tag{3.9}$$

For further use, we also define for any $a = 1, 2$ the number operators $N_a = a_a^\dagger a_a$ satisfying

$$N_a|n\rangle = \theta n|n\rangle \quad \forall |n\rangle \in \mathcal{H}_0. \tag{3.10}$$

To functions on $\mathbb{R}^4$ we associate via the quantization map (3.7) normal ordered operators. On using the number base (3.8) together with (3.9) we may rewrite (3.7) as

$$\hat{\phi} = \sum_{P,Q \in \mathbb{N}^2} \phi_{PQ}|P\rangle\langle Q| \quad \phi_{MN} \in \mathbb{C} \tag{3.11}$$

with $\tilde{\phi}_{PQ}, \phi_{LK}$ related by a change of base. We have indeed

$$|P\rangle = \frac{a_1^{\dagger p_1} a_2^{\dagger p_2}}{[P!\theta^{|P|}]^{1/2}}|0\rangle, \ \ \forall P = (p_1, p_2) \in \mathbb{N}^2, \tag{3.12}$$

with $P! := p_1!p_2!$, $|P| := p_1 + p_2$, and $|0\rangle = |0, 0\rangle$ a Fock vacuum state. This implies

$$\langle z_1, z_2|P\rangle = \overline{\langle P|z_1, z_2\rangle} = e^{-\frac{\bar{z}_1 z_1 + \bar{z}_2 z_2}{2\theta}} \frac{\bar{z}_1^{p_1} \bar{z}_2^{p_2}}{P!\theta^{|P|}}. \tag{3.13}$$

Thus

$$\phi_{LK} = \sum_{q_1=0}^{min\,(l_1,k_1)} \sum_{q_2=0}^{min\,(l_2,k_2)} \tilde{\phi}_{l_2-q_2,k_2-q_2} \frac{\sqrt{L!K!\theta^{|L|+|K|}}}{\theta^{|Q|}Q!}. \tag{3.14}$$

On applying the dequantization map we obtain a function in the noncommutative Voros algebra

$$\phi(z, \bar{z}) = \sum_{PQ} \phi_{PQ} f_{PQ}(z, \bar{z}) \tag{3.15}$$

with

$$f_{PQ}(z, \bar{z}) = \langle z_1, z_2|\hat{f}_{PQ}|z_1, z_2\rangle = \frac{e^{-\frac{\bar{z}_1 z_1 + \bar{z}_2 z_2}{\theta}}}{\sqrt{P!Q!\theta^{|P+Q|}}} \bar{z}_1^{p_1} \bar{z}_2^{p_2} z_1^{q_1} z_2^{q_2} \tag{3.16}$$

and we have introduced the notation $\hat{f}_{PQ} = |P\rangle\langle Q|$. Here we have used (3.13).

The base operators $\hat{f}_{PQ}$ fulfill the following fusion rule, approximation of the identity and trace property respectively given by:

$$\hat{f}_{MN}\hat{f}_{PQ} = \delta_{NP}\hat{f}_{MQ}, \ \mathbf{1} = \sum_{M \in \mathbb{N}^2} |M\rangle\langle M|, \ Tr(\hat{f}_{MN}) = \delta_{MN}, \forall M, N, P, Q \in \mathbb{N}^2. \tag{3.17}$$

These properties descend to the symbol functions of the base $\hat{f}_{MN}$, defining an orthogonal matrix base for $\mathbb{R}^4_\theta$ with a simple rule for the star product

$$f_{MN} \star f_{PQ}(z, \bar{z}) = \langle z_1, z_2|\hat{f}_{MN}\hat{f}_{PQ}|z_1, z_2\rangle = \delta_{NP} f_{MQ}(z, \bar{z}) \tag{3.18}$$

and analogous expressions for the other relations. In particular, on using the decomposition of the identity in terms of coherent states

$$\mathbf{1} = \frac{1}{(\pi\theta)^2} \int d^2 z_1 d^2 z_2 |z_1, z_2\rangle\langle z_1, z_2| \tag{3.19}$$

and the last identity in (3.17), we arrive at

$$\int \mathrm{d}^2 z_1 \mathrm{d}^2 z_2 \, f_{PQ}(z, \bar{z}) = (\pi\theta)^2 \delta_{PQ}. \tag{3.20}$$

The same result can be obtained by direct calculation on using (2.1) and (3.16).

The existence of an orthogonal matrix base for $\mathbb{R}^4_\theta$ allows to rewrite the Voros product in (2.1) as a matrix multiplication. To this we introduce the notation $\Phi := \{\phi_{PK}\}_{P,K \in \mathbb{N}^2}$ for the infinite matrix whose entries are the fields components. We have then

$$\phi \star \psi \, (z, \bar{z}) = \sum_{PQ} \sum_{LK} \phi_{PQ} \psi_{LK} (f_{PQ} \star f_{LK})(z, \bar{z}) = \sum_{PK} (\Phi \cdot \Psi)_{PK} f_{PK}(z, \bar{z}) \tag{3.21}$$

with ($\cdot$) the matrix product. As already noticed, this is a slight modification, adapted to the Voros product, of the matrix base defined in [32] for the Moyal algebra. It was introduced in [40] and already used in [42] in the context of renormalizable scalar field theories on the Voros plane. In the context of quantum field theories approximated with fuzzy geometries the Voros base has been recently used in [43].

3.2 The matrix base of $\mathbb{R}^3_\lambda$

In order to obtain a matrix base in three dimensions, compatible with the product (2.10), we resort to the Schwinger-Jordan realization of the $\mathfrak{su}(2)$ Lie algebra in terms of creation and annihilation operators, which was given in (2.7). The derivation is identical to the one performed in [44] except for the fact that the starting point, the matrix base on $\mathbb{R}^4_\theta$ is here the Voros one.

It is known that $\mathcal{H} = \mathcal{H}_0 \otimes \mathcal{H}_0$ admits the natural decomposition $\mathcal{H} = \bigoplus_{j \in \frac{\mathbb{N}}{2}} \mathcal{V}_j$ where

$$\mathcal{V}_j = span\{|j, m\rangle\}_{-j \le m \le j}, \;\; |j, m\rangle := |j + m\rangle \otimes |j - m\rangle \tag{3.22}$$

is the linear space carrying the irreducible representation of $SU(2)$ with dimension $2j + 1$ and for any $j \in \frac{\mathbb{N}}{2}$, the system $\{|j, m\rangle\}_{-j \le m \le j}$ is orthonormal. From this, it can be realized that another natural base for $\mathbb{R}^4_\theta$ is provided by

$$\{\hat{v}^{j\tilde{j}}_{m\tilde{m}} := |j, m\rangle\langle \tilde{j}, \tilde{m}|\}, \;\; j, \tilde{j} \in \frac{\mathbb{N}}{2}, \;\; -j \le m \le j \, , -\tilde{j} \le \tilde{m} \le \tilde{j} \, . \tag{3.23}$$

The two bases are related as follows. We observe that the eigenvalues of the number operators $\hat{N}_1 = a_1^\dagger a_1$, $\hat{N}_2 = a_2^\dagger a_2$, say p_1, p_2, are related to the eigenvalues of $\hat{\mathbf{X}}^2, \hat{X}_3$, respectively $j(j+1)$ and m, by

$$p_1 + p_2 = 2j \quad p_1 - p_2 = 2m \tag{3.24}$$

with $p_i \in \mathbb{N}$, $j \in \mathbb{N}/2$, $-j \le m \le j$, so to have

$$|p_1 p_2\rangle = |j + m, j - m\rangle \equiv |jm\rangle = \frac{(a_1^\dagger)^{j+m} (a_2^\dagger)^{j-m}}{\sqrt{(j + m)!(j - m)!\theta^{2j}}} |00\rangle \tag{3.25}$$

where $\hat{X}_i, i = 1,..,3$ are the standard angular momentum operators representing the $\mathfrak{su}(2)$ Lie algebra in terms of selfadjoint operators on the Hilbert space $\mathcal{V}_j$ spanned by $|j,m\rangle$. Thus we may identify

$$\hat{f}_{MN} \to \hat{v}^{j\tilde{j}}_{m\tilde{m}}, \tag{3.26}$$

and, for their symbols

$$f_{MN}(z,\bar{z}) \to v^{j\tilde{j}}_{m\tilde{m}}(z,\bar{z}) = \langle z_1, z_2|\hat{v}^{j\tilde{j}}_{m\tilde{m}}|z_1, z_2\rangle \tag{3.27}$$

so to have, for $\phi \in \mathbb{R}^4_\theta$

$$\phi(z_a,\bar{z}_a) = \sum_{j,\tilde{j}\in\mathbb{N}/2} \sum_{m=-j}^{j} \sum_{\tilde{m}=-\tilde{j}}^{\tilde{j}} \phi^{j\tilde{j}}_{m\tilde{m}} v^{j\tilde{j}}_{m\tilde{m}}(z_a,\bar{z}_a) \tag{3.28}$$

We further observe that, for $\phi(z,\bar{z})$ to be in the subalgebra $\mathbb{R}^3_\lambda$ we must impose $j = \tilde{j}$. To this it suffices to compute

$$x_0 \star v^{j\tilde{j}}_{m\tilde{m}}(z,\bar{z}) - v^{j\tilde{j}}_{m\tilde{m}} \star x_0(z,\bar{z}) = \lambda(j-\tilde{j})v^{j\tilde{j}}_{m\tilde{m}} \tag{3.29}$$

with $x_0(z,\bar{z}) = \lambda/(2\theta)\bar{z}_a z_a$ and the $\star$ product defined in (2.10). Then we recall that $\mathbb{R}^3_\lambda$ may be alternatively defined as the $\star$-commutant of x_0. This imposes

$$j = \tilde{j} \tag{3.30}$$

We have then

$$\phi(x_i,x_0) = \sum_j \sum_{m,\tilde{m}=-j}^{j} \phi^j_{m\tilde{m}} v^j_{m\tilde{m}} \tag{3.31}$$

with

$$v^j_{m\tilde{m}} := v^{jj}_{m\tilde{m}} = e^{-\frac{\bar{z}_a z_a}{\theta}} \frac{\bar{z}_1^{j+m} z_1^{j+\tilde{m}} \bar{z}_2^{j-m} z_2^{j-\tilde{m}}}{\sqrt{(j+m)!(j-m)!(j+\tilde{m})!(j-\tilde{m})!\theta^{4j}}} \tag{3.32}$$

and we recall its expression in terms of the dequantization map

$$v^j_{m\tilde{m}}(z,\bar{z}) = \langle z_1, z_2|j\, m\rangle\langle j\, \tilde{m}|z_1, z_2\rangle. \tag{3.33}$$

As for the normalization we have

$$\int \mathrm{d}^2 z_1 \mathrm{d}^2 z_2 \; v^j_{m\tilde{m}}(z,\bar{z}) = \pi^2\theta^2\delta_{m\tilde{m}}. \tag{3.34}$$

Let us notice that the base elements $v^j_{m\tilde{m}}(z,\bar{z})$ can be reexpressed solely in terms of the coordinate functions x_i, x_0 although the expression is not unique. A possible choice is

$$v^j_{m\tilde{m}}(x_i,x_0) = \kappa\; e^{-2\lambda x_0} \frac{(x_0+x_3)^{j+m}(x_0-x_3)^{j-\tilde{m}}\;(x_1-ix_2)^{\tilde{m}-m}}{\sqrt{(j+m)!(j-m)!(j+\tilde{m})!(j-\tilde{m})!\lambda^{2j}}}. \tag{3.35}$$

The star product acquires the simple form

$$v^j_{m\tilde{m}} \star v^{\tilde{j}}_{n\tilde{n}} = \delta^{j\tilde{j}}\delta_{\tilde{m}n} v^j_{m\tilde{n}} \tag{3.36}$$

which implies the orthogonality property

$$\int v^j_{m\tilde{m}} \star v^{\tilde{j}}_{n\tilde{n}} = \pi^2\theta^2\delta^{j\tilde{j}}\delta_{\tilde{m}n}\delta_{m\tilde{n}}. \tag{3.37}$$

These properties may be either directly verified or derived from the dequantization map starting from the operator relations

$$\hat{v}^{j_1}_{m_1,m_2}\hat{v}^{j_2}_{n_1,n_2} = \delta_{j_1 j_2}\delta_{m_2 n_1}\hat{v}^{j_1}_{m_1,n_2},\ (\hat{v}^{j}_{m_1,m_2})^\dagger = \hat{v}^{j}_{m_2,m_1}, \tag{3.38}$$

$$\langle \hat{v}^{j_1}_{m_1,m_2}, \hat{v}^{j_2}_{n_1,n_2}\rangle = \delta_{j_1 j_2}\delta_{m_1 n_1}\delta_{m_2 n_2} \quad \mathbf{1} = \sum_{j\in\frac{\mathbb{N}}{2}}\sum_{m=-j}^{j}\hat{v}^{j}_{mm},\ \ \mathrm{Tr}\,(\hat{v}^{j}_{m_1,m_2}) = \delta_{m_1 m_2} \tag{3.39}$$

where $\langle \hat{v}^{j_1}_{m_1,m_2}, \hat{v}^{j_2}_{n_1,n_2}\rangle := \ \mathrm{Tr}\,(\hat{v}^{j_1}_{m_1,m_2})^\dagger \hat{v}^{j_2}_{n_1,n_2}$ is the scalar product.

Notice that, for any $j \in \frac{\mathbb{N}}{2}$, the set $\{\hat{v}^{j}_{m_1,m_2}\}$, $-j \le m_1, m_2 \le j$ of $(2j+1)^2$ linear maps $v^{j}_{m_1,m_2} : \mathcal{V}^j \to \mathcal{V}^j$ simply describes the canonical base of the algebra of endomorphisms of $\mathcal{V}^j$, orthonormal with respect to the scalar product introduced above. From this it follows that the direct sum decomposition

$$\mathbb{R}^3_\lambda \simeq \bigoplus_{j\in\frac{\mathbb{N}}{2}} End(\mathcal{V}^j) \simeq \bigoplus_{j\in\frac{\mathbb{N}}{2}} \mathbb{S}^j \tag{3.40}$$

holds true, where $End(\mathcal{V}^j)$ denotes the algebra of endomorphisms of $\mathcal{V}^j$, $\forall j \in \frac{\mathbb{N}}{2}$, which actually describe the so-called fuzzy spheres of different radii, $\mathbb{S}^j$.

The star product in $\mathbb{R}^3_\lambda$ becomes a block-diagonal matrix product

$$\begin{aligned}\phi\star\psi(x_i,x_0) &= \sum \phi^j_{m_1\tilde{m}_1}\psi^j_{m_2\tilde{m}_2} v^j_{m_1\tilde{m}_1}\star v^j_{m_2\tilde{m}_2} = \sum \phi^j_{m_1\tilde{m}_1}\psi^j_{m_2\tilde{m}_2} v^j_{m_1\tilde{m}_2}\delta_{\tilde{m}_1 m_2} \\ &= \sum_{j,m_1,\tilde{m}_2} (\Phi^j\cdot\Psi^j)_{m_1\tilde{m}_2} v^j_{m_1\tilde{m}_2}\end{aligned} \tag{3.41}$$

where the infinite matrix Φ gets rearranged into a block-diagonal form, each block being the $(2j+1)\times(2j+1)$ matrix $\Phi^j = \{\phi^j_{mn}\}$, $-j \le m, n \le j$. The integral may be defined through the pullback to $\mathbb{R}^4_\theta$

$$\int_{\mathbb{R}^3_\lambda} \phi := \frac{\kappa^3}{\pi^2\theta^2}\int_{\mathbb{R}^4_\theta} \pi^\star(\phi) = \kappa^3 \sum_j \ \mathrm{Tr}_{\,j}\Phi^j \tag{3.42}$$

with $\mathrm{Tr}_{\,j}$ the trace in the $(2j+1)\times(2j+1)$ subspace[¶]. We have also

$$\int_{\mathbb{R}^3_\lambda} \phi\star\psi := \frac{\kappa^3}{\pi^2\theta^2}\int_{\mathbb{R}^4_\theta} \pi^\star(\phi)\star\pi^*(\psi) = \kappa^3\sum_j \ \mathrm{Tr}_{\,j}\Phi^j\Psi^j. \tag{3.43}$$

4 The scalar actions

In this section we consider a family of scalar field theories on $\mathbb{R}^3_\lambda$ indexed by two real parameters α, β. We assume the fields $\phi \in R^3_\lambda$ to be real. Upon rewriting the action in the matrix base we perform one loop calculations and discuss the divergences. Some comments on the renormalization of the theory are given at the end.

[¶] If we were to perform our analysis in the coordinate base, without recurring to the matrix base, we should use a differential calculus adapted to $\mathbb{R}^3_\lambda$ as the one introduced in [45] .

4.1 General properties

Let

$$S[\phi] = \int \phi \star (\Delta + \mu^2)\phi + \frac{g}{4!}\phi \star \phi \star \phi \star \phi \tag{4.1}$$

where Δ is the Laplacian defined as

$$\Delta\phi = \alpha(D_i^2)\phi + \frac{\beta}{\kappa^4} x_0 \star x_0 \star \phi = \frac{1}{\kappa^4}\big(\alpha[x_i, [x_i, \phi]_\star]_\star + \beta x_0 \star x_0 \star \phi\big) \tag{4.2}$$

and $D_i = \kappa^{-2}[x_i, \ \cdot \ ]_\star$, $i = 1, .., 3$ are inner derivations of $\mathbb{R}^3_\lambda$. The mass dimensions are $[\phi] = \frac{1}{2}$, $[g] = 1$, $[D_i] = 1$. α and β are dimensionless parameters.

The second term in the Laplacian has been added in order to reproduce the radial part of the usual Laplacian on the commutative $\mathbb{R}^3$. From (2.10) we have indeed

$$[x_i, \phi]_\star = -i\lambda\epsilon_{ijk}x_j\partial_k\phi \tag{4.3}$$

so that the first term, that is $[x_i, [x_i, \phi]_\star]\star$ reproduces the angular part of the Laplacian on $\mathbb{R}^3$, whereas

$$x_0 \star \phi = x_0\phi + \frac{\lambda}{2}x_i\partial_i\phi \tag{4.4}$$

contains the dilation operator. Therefore, the highest derivatives term of the Laplacian defined in (4.2) can be made into the ordinary Laplacian on $\mathbb{R}^3$ for the parameters α and β appropriately chosen.

To rewrite the action in the matrix base $\{v^j_{m\tilde{m}}\}$ we first express the coordinate functions in such a base. On using the expression of the generators in terms of $\bar{z}_a, z_a$, (2.7) and the base transformations (3.14) we find

$$\begin{aligned}
x_+ &= \frac{\lambda}{\theta}\bar{z}_1 z_2 = \lambda\sum_{j,m}\sqrt{(j+m)(j-m+1)}v^j_{m\,m-1} && (4.5)\\
x_- &= \frac{\lambda}{\theta}\bar{z}_2 z_1 = \lambda\sum_{j,m}\sqrt{(j-m)(j+m+1)}v^j_{m\,m+1} && (4.6)\\
x_3 &= \frac{\lambda}{2\theta}(\bar{z}_1 z_1 - \bar{z}_2 z_2) = \lambda\sum_{j,m} m v^j_{m\,m} && (4.7)\\
x_0 &= \frac{\lambda}{2\theta}(\bar{z}_1 z_1 + \bar{z}_2 z_2) = \lambda\sum_{j,m} j v^j_{m\,m} && (4.8)
\end{aligned}$$

were we have introduced

$$x_\pm := x_1 \pm i x_2. \tag{4.9}$$

Thus we compute

$$\begin{array}{ll}
x_+ \star v^j_{m\tilde{m}} = \lambda\sqrt{(j+m+1)(j-m)}v^j_{m+1\,\tilde{m}} & v^j_{m\tilde{m}} \star x_+ = \lambda\sqrt{(j-\tilde{m}+1)(j+\tilde{m})}v^j_{m\,\tilde{m}-1}\\
x_- \star v^j_{m\tilde{m}} = \lambda\sqrt{(j-m+1)(j+m)}v^j_{m-1\,\tilde{m}} & v^j_{m\tilde{m}} \star x_- = \lambda\sqrt{(j+\tilde{m}+1)(j-\tilde{m})}v^j_{m\,\tilde{m}+1}\\
x_3 \star v^j_{m\tilde{m}} = \lambda\, m\, v^j_{m\tilde{m}} & v^j_{m\tilde{m}} \star x_3 = \lambda\,\tilde{m}\, v^j_{m\tilde{m}}\\
x_0 \star v^j_{m\tilde{m}} = \lambda\, j\, v^j_{m\tilde{m}} & v^j_{m\tilde{m}} \star x_0 = \lambda\, j\, v^j_{m\tilde{m}}
\end{array} \tag{4.10}$$

which yield

$$\begin{aligned}
[x_+,[x_-,v^j_{m\tilde m}]_\star]_\star &= \lambda^2\Big\{\Big((j+m)(j-m+1)+(j+\tilde m+1)(j-\tilde m)\Big)v^j_{m\tilde m}+\\
&- \sqrt{(j+m)(j-m+1)(j+\tilde m)(j-\tilde m+1)}v^j_{m-1\tilde m-1}+\\
&- \sqrt{(j+m+1)(j-m)(j+\tilde m+1)(j-\tilde m)}v^j_{m+1\tilde m+1}\Big\}
\end{aligned}\tag{4.11}$$

$$\begin{aligned}
[x_-,[x_+,v^j_{m\tilde m}]_\star]_\star &= \lambda^2\Big\{\Big((j+m+1)(j-m)+(j+\tilde m)(j-\tilde m+1)\Big)v^j_{m\tilde m}+\\
&- \sqrt{(j+m)(j-m+1)(j+\tilde m)(j-\tilde m+1)}v^j_{m-1\tilde m-1}+\\
&- \sqrt{(j+m+1)(j-m)(j+\tilde m+1)(j-\tilde m)}v^j_{m+1\tilde m+1}\Big\}
\end{aligned}\tag{4.12}$$

$$\begin{aligned}
[x_3,[x_3,v^j_{m\tilde m}]_\star]_\star &= \lambda^2(m-\tilde m)^2v^j_{m\tilde m}\\
x_0\star x_0\star v^j_{m\tilde m} &= \lambda^2 j^2 v^j_{m\tilde m}
\end{aligned}\tag{4.13}$$

These relations allow to compute

$$\Delta(\alpha,\beta)v^j_{m\tilde m}=\frac{\alpha}{\kappa^4}\Big(\frac{1}{2}([x_+,[x_-,v^j_{m\tilde m}]_\star]_\star+[x_-,[x_+,v^j_{m\tilde m}]_\star]_\star)+x_3,[x_3,v^j_{m\tilde m}]_\star]_\star\Big)+\frac{\beta}{\kappa^4}x_0\star x_0\star v^j_{m\tilde m}\tag{4.14}$$

On using the expansion of the fields in the matrix base and the multiplication rule for the base elements, already introduced in the previous section, respectively

$$\phi=\sum_{j,m\tilde m}\phi^j_{m\tilde m}v^j_{m\tilde m}\tag{4.15}$$

and

$$v^j_{m\tilde m}\star v^j_{n\tilde n}=\delta_{\tilde m n}v^j_{m\tilde n}\tag{4.16}$$

we obtain the action in (4.1) as a matrix model action

$$\begin{aligned}
S[\phi] &= \kappa^3\big\{\sum\phi^j_{m_1\tilde m_1}\big(\Delta(\alpha,\beta)+\mu^2\mathbf{1}\big)^j_{m_1\tilde m_1;m_2\tilde m_2}\phi^j_{m_2\tilde m_2}+\frac{g}{4!}\sum\phi^j_{mn}\phi^j_{np}\phi^j_{pq}\phi^j_{qm}\big\}\\
&= \kappa^3\big\{\,\mathrm{Tr}\,(\Phi(\Delta(\alpha,\beta)+\mu^2\mathbf{1})\Phi)+\frac{g}{4!}\,\mathrm{Tr}\,(\Phi\Phi\Phi\Phi)\big\}
\end{aligned}\tag{4.17}$$

where sums are understood over all the indices and $\mathrm{Tr}\,:=\sum_j\,\mathrm{Tr}\,_j$. The matrix elements of the identity operator are

$$\mathbf{1}^j_{m_1\tilde m_1 m_2\tilde m_2}=\delta_{\tilde m_1 m_2}\delta_{m_1\tilde m_2}\tag{4.18}$$

The kinetic operator may be verified to be

$$\begin{aligned}
(\Delta(\alpha,\beta)+\mu^2\mathbf{1})^j_{m_1\tilde m_1;m_2\tilde m_2} &:= \frac{1}{\pi^2\theta^2}\int v^j_{m_1\tilde m_1}\star(\Delta(\alpha,\beta)+\mu^2\mathbf{1})v^j_{m_2\tilde m_2}\\
&= \frac{\lambda^2}{\kappa^4}\big\{\delta_{\tilde m_1 m_2}\delta_{m_1\tilde m_2}D^j_{m_2\tilde m_2}-\delta_{\tilde m_1,m_2+1}\delta_{m_1,\tilde m_2+1}B^j_{m_2,\tilde m_2}\\
&- \delta_{\tilde m_1,m_2-1}\delta_{m_1,\tilde m_2-1}H^j_{m_2,\tilde m_2}\,\big\}
\end{aligned}\tag{4.19}$$

with

$$D^j_{m_2\tilde m_2} = [(2\alpha+\beta)j^2+2\alpha(j_2-m_2\tilde m_2)]+\lambda^2\mu^2\tag{4.20}$$

$$B^j_{m_2\tilde m_2} = \alpha\sqrt{(j+m_2+1)(j-m_2)(j+\tilde m_2+1)(j-\tilde m_2)}\tag{4.21}$$

$$H^j_{m_2\tilde m_2} = \alpha\sqrt{(j+m_2)(j-m_2+1)(j+\tilde m_2)(j-\tilde m_2+1)}.\tag{4.22}$$

At this stage, some comments are in order.

i) The use of the matrix base for $\mathbb{R}^3_\lambda$ yields an interaction term which is diagonal (i.e a simple trace of product of matrices built from the coefficients of the fields expansion) whereas the kinetic term is not diagonal. Had we used the expansion of ϕ in the so called fuzzy harmonics base (Y^j_{lk}), $j \in \frac{\mathbb{N}}{2}$, $l \in \mathbb{N}$, $0 \leq l \leq 2j$, $-l \leq k \leq l$ (see below), then we would have obtained a diagonal kinetic term with a complicated interaction term. Notice that this remark holds for any polynomial interaction term. We will come back to this point in a while.

ii) We observe that the action (4.17) is expressed as an infinite sum of contributions, namely $S[\Phi] = \sum_{j\in\frac{\mathbb{N}}{2}} S^{(j)}[\Phi]$, where the expression for $S^{(j)}$ can be read off from (4.17) and describes a scalar action on the fuzzy sphere $\mathbb{S}^j \simeq End(\mathcal{V}^j)$.

A lot of information about the short and long distance behaviour of a matrix model with diagonal interaction term, regarding renormalization properties, is encoded into the propagator. The computation of this latter amounts to the determination of the inverse of the kinetic operator in Eq. (4.17) operator which, because of the remark ii) above, is expressible into a block diagonal form. Explicitly

$$S_{Kin}[\Phi] = \kappa^3 \sum_j \sum_{m,\tilde{m}} \phi^j_{m_1\tilde{m}_1}(\Delta + \mu^2\mathbf{1})^j_{m_1\tilde{m}_1;m_2\tilde{m}_2}\phi^j_{m_2\tilde{m}_2}, \tag{4.23}$$

with $\Delta^j_{m_1\tilde{m}_1;m_2\tilde{m}_2}$ defined in (4.19). Since the mass term is diagonal, let us put it to zero for the moment. We shall restore it at the end. One has the following "law of indices conservation"

$$\Delta^j_{mn;kl} = 0 \iff m + k \neq n + l \tag{4.24}$$

We denote by $P^j_{mn;kl}(\alpha,\beta)$ the inverse of $\Delta^j_{mn;kl}(\alpha,\beta)$ which is defined by

$$\sum_{k,l=-j}^{j} \Delta^j_{mn;lk}P^j_{lk;rs} = \delta_{ms}\delta_{nr}, \quad \sum_{m,n=-j}^{j} P^j_{rs;mn}\Delta^j_{mn;kl} = \delta_{rl}\delta_{sk}, \tag{4.25}$$

for which the law of indices conservation still holds true as

$$P^j_{mn;kl} = 0 \iff m + k \neq n + l. \tag{4.26}$$

To determine $P^j_{mn;kl}$ one has to diagonalize $\Delta^j_{mn;kl}$. This can be done by brute force, as in [16] for the case of noncommutative scalar theories on the Moyal spaces $\mathbb{R}^{2n}_\theta$, by first exploiting the consequence of (4.24), (4.26) to turn the multi-indices quantities $\Delta^j_{mn;kl}$ and $P^j_{mn;kl}$ (4.25) to two sets (indexed by, say, $m-n = l-k$, see (4.24), (4.26)) of matrices with two indices, then by looking for a set of unitary transformations diagonalising the set of matrices stemming from $\Delta^j_{mn;kl}(\alpha,\beta)$. Then, each of these unitary transformations is found to be a solution of a 3-terms recursive equation defining a particular class of orthogonal polynomials. In the case of the scalar noncommutative field theories considered in [16], the above recursive equations are solved by a specific family of Meixner polynomials [50]. Having diagonalized the kinetic operator, the expression for the propagator follows. In the present case, a similar direct calculation can be performed and gives rise after some calculations to a family of 3-term recursive equations, solved by a particular family of polynomials, the so called dual Hahn polynomials [50]. Details of this derivation are given in Appendix. However the whole computation of the propagator can be considerably simplified observing that, as already mentioned, we already know an alternative orthogonal base for $End(\mathcal{V}^j)$ where the kinetic part of the action can be shown to be diagonal. These are the so called fuzzy spherical harmonics. We show in the appendix that the two methods are equivalent, and we exhibit the proportionality relation between dual Hahn polynomials and the fuzzy spherical harmonics.

4.2 The kinetic action in the fuzzy spherical harmonics base

It is well known and largely exploited in the literature on the fuzzy sphere, that $End(\mathcal{V}^j)$ is spanned by the so called Fuzzy Spherical Harmonics Operators, or, up to normalization factors, irreducible tensor operators. We shall indicate them as

$$\hat{Y}^j_{lk} \in End(\mathcal{V}^j), \quad l \in \mathbb{N}, \quad 0 \le l \le 2j, \quad -l \le k \le l, \tag{4.27}$$

whereas the unhatted objects Y^j_{lk} are their symbols and are sometimes referred to as fuzzy spherical harmonics with no other specification (notice however that the functional form of the symbols does depend on the dequantization map that has been chosen). Concerning the definition and normalization of the fuzzy spherical harmonics operators, we use the following conventions [46]. We set

$$J_\pm = \frac{\hat{x}_\pm}{\lambda}. \tag{4.28}$$

We have, for $l = m$,

$$\hat{Y}^j_{ll} := (-1)^l \frac{\sqrt{2j+1}}{l!} \frac{\sqrt{(2l+1)!(2j-l)!}}{(2j+l+1)!} (J_+)^l \tag{4.29}$$

while the others are defined recursively through the action of J_-

$$\hat{Y}^j_{lk} := [(l+k+1)(l-k)]^{-\frac{1}{2}} [J_-, \hat{Y}^j_{l,k+1}], \tag{4.30}$$

and satisfy

$$(\hat{Y}^j_{lk})^\dagger = (-1)^{k-2j} \hat{Y}^j_{l,-k}, \; \langle \hat{Y}^j_{l_1k_1}, \hat{Y}^j_{l_2k_2} \rangle = \operatorname{Tr}((\hat{Y}^j_{l_1k_1})^\dagger \hat{Y}^j_{l_2k_2}) = (2j+1)\delta_{l_1l_2}\delta_{k_1k_2}. \tag{4.31}$$

The symbols are defined through the dequantization map (3.4)

$$Y^j_{lk} := \langle z | \hat{Y}^j_{lk} | z \rangle. \tag{4.32}$$

From (4.30), (4.31) and the Lie algebra relation $[J_+, J_-] = 2J_3$ it is straightforward to check the usual properties

$$\begin{aligned}
[J_-, \hat{Y}^j_{lk}] &= \sqrt{(l+k)(l-k+1)} \hat{Y}^j_{l\,k-1} && (4.33)\\
[J_+, \hat{Y}^j_{lk}] &= \sqrt{(l-k)(l+k+1)} \hat{Y}^j_{l\,k+1} && (4.34)\\
[J_3, \hat{Y}^j_{lk}] &= k\, \hat{Y}^j_{lk} && (4.35)\\
[J_i, [J_i, \hat{Y}^j_{lk}]] &= l(l+1) \hat{Y}^j_{lk} && (4.36)
\end{aligned}$$

which imply for the symbols

$$\begin{aligned}
[x_-, Y^j_{lk}]_\star &= \lambda < z|[J_-, \hat{Y}^j_{lk}]|z >= \lambda \sqrt{(l+k)(l-k+1)} Y^j_{l\,k-1} \\
[x_+, Y^j_{lk}]_\star &= \lambda < z|[J_+, \hat{Y}^j_{lk}]|z >= \lambda \sqrt{(l-k)(l+k+1)} Y^j_{l\,k+1} \\
[x_3, Y^j_{lk}]_\star &= \lambda < z|[J_3, \hat{Y}^j_{lk}]|z >= \lambda\, k\, Y^j_{lk}
\end{aligned} \tag{4.37}$$

and in particular

$$[x_i, [x_i, Y^j_{lk}]_\star]_\star = \lambda^2 < z|[J_i, [J_i, \hat{Y}^j_{lk}]]|z >= \lambda^2\, l(l+1)\, Y^j_{lk}. \tag{4.38}$$

In order to evaluate the action of the full Laplacian (4.2) on the fuzzy spherical harmonics we need to compute $x_0 \star Y^j_{lk}$. To this we express the fuzzy spherical harmonics in the canonical base $v^j_{m\tilde{m}}$

$$Y^j_{lk} = \sum_{-j\leq m,\tilde{m}\leq j} (Y^j_{lk})_{m\tilde{m}} v^j_{m\tilde{m}}, \tag{4.39}$$

where the coefficients are given in terms of Clebsch-Gordan coefficients by [46]

$$(Y^j_{lk})_{m\tilde{m}} = \langle \hat{v}^j_{m\tilde{m}} | \hat{Y}^j_{lk} \rangle = \sqrt{2j+1}(-1)^{j-\tilde{m}} \left(\begin{array}{cc|c} j & j & l \\ m & -\tilde{m} & k \end{array} \right), \; -j \leq m, \tilde{m} \leq j\,, \tag{4.40}$$

$$(Y^{j\,\dagger}_{lk})_{m\tilde{m}} = (-1)^{-2j} (Y^j_{lk})_{\tilde{m}m}. \tag{4.41}$$

On using Eq. (4.39), and the orthogonality relation of Clebsch-Gordan coefficients

$$\sum_{m\tilde{m}} \left(\begin{array}{cc|c} j & j & l_1 \\ m & \tilde{m} & k_1 \end{array} \right) \left(\begin{array}{cc|c} j & j & l_2 \\ m & \tilde{m} & k_2 \end{array} \right) = \delta_{l_1 l_2} \delta_{k_1 k_2} \tag{4.42}$$

together with the star product (3.37) it is straightforward to check that

$$\int {Y^j}_{l_1k_1} \star Y^j_{l_2k_2} = \kappa^3 (-1)^{k_1+2j} (2j+1) \delta_{l_1l_2} \delta_{-k_1k_2} \tag{4.43}$$

$$\int {Y^j}^\dagger_{l_1k_1} \star Y^j_{l_2k_2} = \kappa^3 (2j+1) \delta_{l_1l_2} \delta_{k_1k_2} \tag{4.44}$$

in accordance with the second of relations (4.31). Eq. (4.39), and last of Eqs. (4.10) imply

$$x_0 \star Y^j_{lk} = \sum_{-j\leq m,\tilde{m}\leq j} (Y^j_{lk})_{m\tilde{m}} \; x_0 \star v^j_{m\tilde{m}} = \lambda j Y^j_{lk}. \tag{4.45}$$

Thus, from the definition of the Laplacian (4.2) and Eq. (4.38), Eq. (4.45), we verify that in the fuzzy spherical harmonics base the whole kinetic term is diagonal,

$$\Delta(\alpha,\beta) Y^j_{lk} = \frac{\lambda^2}{\kappa^4} \left(\alpha l(l+1) + \beta j^2 \right) Y^j_{lk} \quad j \in \frac{\mathbb{N}}{2}, \; 0 \leq l \leq 2j, \; l \in \mathbb{N}, \; -l \leq k \leq l \tag{4.46}$$

with eigenvalues

$$\gamma(j,l;\alpha,\beta) := \alpha l(l+1) + \beta j^2. \tag{4.47}$$

Note that (4.46), (4.47), with our choice for the dimensionality of the parameters λ, κ, single out a natural choice for the UV and IR regimes, which correspond respectively to large or small values of $\gamma(j,l;\alpha,\beta)$. We can expand the fields $\phi \in \mathbb{R}^3_\lambda$ in the fuzzy harmonics base

$$\phi = \sum_{j\in\frac{\mathbb{N}}{2}} \sum_{l=0}^{2j} \sum_{k=-l}^{l} \varphi^j_{lk} Y^j_{lk}, \tag{4.48}$$

and comparing with their expression in the canonical base, (4.15), we readily obtain

$$\phi^j_{m\tilde{m}} = \sum_{l=0}^{2j} \sum_{k=-l}^{l} (Y^j_{lc})_{m\tilde{m}} \varphi^j_{lk} = \sum_{l=0}^{2j} \sum_{k=-l}^{l} \sqrt{2j+1} (-1)^{j-m_2} \left(\begin{array}{cc|c} j & j & l \\ m & -\tilde{m} & k \end{array} \right) \varphi^j_{lk}, \tag{4.49}$$

which relates the propagating degree of freedom φ^j_{lk}, for which the kinetic term of the action is diagonal, to the interacting degree of freedom $\phi^j_{m\tilde{m}}$ for which the interaction term is diagonal.

From (4.48) and the properties of the fuzzy harmonics we derive, for a Hermitian field ϕ,

$$\phi^\dagger = \phi \;\Rightarrow\; \varphi^j_{lk} = (-1)^{-k-2j}\varphi^{j*}_{l,-k}, \qquad j \in \frac{\mathbb{N}}{2},\; l \in \mathbb{N},\; 0 \le l \le 2j,\; -l \le k \le l. \tag{4.50}$$

Therefore, upon restoring the mass term, we can compute the kinetic action in the fuzzy harmonics base

$$\begin{aligned} \int \phi \star (\Delta + \mu^2)\phi &= \frac{\lambda^2}{\kappa^4}\sum \varphi^j_{l_1k_1}\varphi^j_{l_2k_2}\left(\gamma(j,l_2;\alpha,\beta) + \frac{\kappa^4}{\lambda^2}\mu^2\right)\int Y^j_{l_1k_1} \star Y^j_{l_2k_2} && (4.51)\\ &= \frac{\lambda^2}{\kappa^4}\sum \varphi^{j*}_{l_1k_1}\varphi^j_{l_2k_2}\left(\gamma(j,l_2;\alpha,\beta) + \frac{\kappa^4}{\lambda^2}\mu^2\right)\int {Y^j}^\dagger_{l_1k_1} \star Y^j_{l_2k_2} \\ &= \frac{\lambda^2}{\kappa}\sum |\varphi^j_{lk}|^2(2j+1)(\gamma(j,l;\alpha,\beta) + \frac{\kappa^4}{\lambda^2}\mu^2) && (4.52) \end{aligned}$$

which is positive for $\alpha, \beta \geq 0$. We define for further convenience

$$(\Delta_{diag})^j_{l_1k_1l_2k_2} = \frac{1}{\lambda^3}\int Y^j_{l_1k_1} \star \Delta(\alpha,\beta)Y^j_{l_2k_2} = \frac{1}{\lambda^2}(-1)^{k_1+2j}(2j+1)\gamma(j,l_2;\alpha,\beta)\delta_{l_1l_2}\delta_{-k_1k_2}. \tag{4.53}$$

4.3 The propagator

We can now state the following Lemma

Lemma 1 *Let $\mathbb{R}^3_\lambda$ be the noncommutative algebra defined in* (3.40)*, with canonical base $\{v^j_{m\tilde{m}}\}$, $j \in \frac{\mathbb{N}}{2}$, $-j \le m, \tilde{m} \le j$, together with the fuzzy spherical harmonics base $\{Y^j_{lk}\}$, $j \in \frac{\mathbb{N}}{2}$, $0 \le l \le 2j$, $-j \le m \le j$. The inverse of the kinetic operator, in the canonical base,*

$$\big(\Delta(\alpha,\beta) + \mu^2\mathbf{1}\big)^j_{m_1\tilde{m}_1;m_2\tilde{m}_2} = \frac{1}{\pi^2\theta^2}\int v^j_{m_1\tilde{m}_1} \star \big(\Delta(\alpha,\beta) + \mu^2\mathbf{1}\big)v^j_{m_2\tilde{m}_2} \tag{4.54}$$

is given by

$$(P(\alpha,\beta))^j_{p_1,\tilde{p}_1;p_2\tilde{p}_2} = \sum_{l=0}^{2j}\sum_{k=-l}^{l}\int_0^\infty dt\; e^{-t(2j+1)(\frac{\lambda^2}{\kappa^4}\gamma(j,l;\alpha,\beta)+\mu^2)}({Y^j_{lk}}^\dagger)_{p_1\tilde{p}_1}(Y^j_{lk})_{p_2\tilde{p}_2}, \tag{4.55}$$

where $\gamma(j,l;\alpha,\beta)$, the eigenvalues of the Laplacian operator, have been given in (4.47).

Proof It is based on the so called Schwinger parametrization. For each positive operator A we can write

$$\frac{1}{A} = \int_0^\infty dt\, e^{-tA} \tag{4.56}$$

This applies to the matrix elements of the kinetic operator in the diagonal (propagation) base

$$\begin{aligned} &\left[(\Delta_{diag} + \mu^2\mathbf{1})^{-1}\right]^j_{l_1k_1l_2k_2} \\ &\quad = \int dt\, e^{-t(\Delta_{diag}+\mu^2\mathbf{1})^j_{l_1k_1l_2k_2}} = (-1)^{k_1+2j}\int dte^{-t(2j+1)(\frac{\lambda^2}{\kappa^4}\gamma(j,l_2;\alpha\beta)+\mu^2)}\delta_{l_1l_2}\delta_{-k_1k_2} \end{aligned} \tag{4.57}$$

Let us perform a change of base from the diagonal to the interaction (canonical) base. We have

$$\begin{aligned}\int \phi \star \Delta\phi &= \kappa^3 \sum \phi^j_{m_1\tilde{m}_1} \Delta^j_{m_1\tilde{m}_1 m_2\tilde{m}_2} \phi^j_{m_2\tilde{m}_2} \\ &= \kappa^3 \sum \varphi^j_{l_1k_1} (Y^j_{l_1k_1})_{m_1\tilde{m}_1} \Delta^j_{m_1\tilde{m}_1 m_2\tilde{m}_2} \varphi^j_{l_2k_2} (Y^j_{l_2k_2})_{m_2\tilde{m}_2} \end{aligned} \tag{4.58}$$

By comparing with the expression in the diagonal base (4.51) we obtain

$$(\Delta^j_{diag})_{l_1k_1l_2k_2} = (Y^j_{l_1k_1})_{m_1\tilde{m}_1} \Delta^j_{m_1\tilde{m}_1 m_2\tilde{m}_2} (Y^j_{l_2k_2})_{m_2\tilde{m}_2} \tag{4.59}$$

with inverse transformation

$$\Delta^j_{m_1\tilde{m}_1 m_2\tilde{m}_2} = \frac{1}{(2j+1)^2} (Y^j_{l_1k_1})_{m_1\tilde{m}_1} (\Delta^j_{diag})_{l_1k_1l_2k_2} (Y^j_{l_2k_2})_{m_2\tilde{m}_2} \tag{4.60}$$

The (massless) propagator is then

$$[\Delta^j)^{-1}]_{m_1\tilde{m}_1 m_2\tilde{m}_2} = (Y^j_{l_1k_1})_{m_1\tilde{m}_1} [(\Delta^j_{diag})^{-1}]_{l_1k_1l_2k_2} (Y^j_{l_2k_2})_{m_2\tilde{m}_2} \tag{4.61}$$

On replacing the expression for the diagonal inverse (4.57) and on using the first of Eqs. (4.31) we arrive at

$$\begin{aligned}[\Delta + \mu^2 \mathbf{1})^{-1}]^j_{m_1\tilde{m}_1 m_2\tilde{m}_2} &= (-1)^{-k+2j} \sum_{l=0}^{2j} \sum_{k=-l}^{l} \int dt e^{-t(2j+1)\left(\frac{\lambda^2}{\kappa^4}\gamma(j,l;\alpha,\beta)+\mu^2\right)} (Y^j_{l-k})_{m_1\tilde{m}_1} (Y^j_{lk})_{m_2\tilde{m}_2} \\ &= \sum_{l=0}^{2j} \sum_{k=-l}^{l} \int dt e^{-t(2j+1)\left(\frac{\lambda^2}{\kappa^4}\gamma(j,l;\alpha,\beta)+\mu^2\right)} (Y^{j\dagger}_{lk})_{m_1\tilde{m}_1} (Y^j_{lk})_{m_2\tilde{m}_2} \end{aligned} \tag{4.62}$$

which completes the proof. The result can be verified directly, by using the orthogonality properties of the fuzzy harmonics. ■

5 One-loop calculations

Once we have established the form of the propagator in the matrix base,

$$P^j_{m_1\tilde{m}_1 m_2\tilde{m}_2} = \sum_{l=0}^{2j} \sum_{k=-l}^{l} \int dt e^{-t(2j+1)\left(\frac{\lambda^2}{\kappa^4}\gamma(j,l;\alpha,\beta)+\mu^2\right)} (Y^{j\dagger}_{lk})_{m_1\tilde{m}_1} (Y^j_{lk})_{m_2\tilde{m}_2} \tag{5.1}$$

and of the vertex

$$V^{j_1j_2j_3j_4}_{p_1\tilde{p}_1;p_2\tilde{p}_2;p_3\tilde{p}_3;p_4\tilde{p}_4} = \frac{g}{4!} \delta^{j_1j_2j_3j_4} \delta_{\tilde{p}_1p_2} \delta_{\tilde{p}_2p_3} \delta_{\tilde{p}_3p_4} \delta_{\tilde{p}_4p_1} \tag{5.2}$$

the computation of Feynman graphs (ribbon graphs) of every order is fairly easy: it is just a matter of gluing together the appropriate number of propagators, which are represented by a double-line while contracting them with the diagonal vertex (5.2). We explicitly compute one-loop planar and non planar contributions to the 2-point correlation function.



$$= P_{m_1\tilde m_1;m_2\tilde m_2}$$

Figure 1: The propagator



Figure 2: The vertex

5.1 Planar two-point Green function

The planar diagram contributing to the 2-point correlation function is depicted on Fig. 3. Its amplitude is given by

$$\begin{aligned}\mathcal{A}_{p_1\tilde p_1;p_2\tilde p_2} &= \sum_{j=0}^{\infty}\sum_{p_3\tilde p_3 p_4\tilde p_4} V^j_{p_1\tilde p_1;p_2\tilde p_2;p_3\tilde p_3;p_4\tilde p_4} P^j_{\tilde p_4 p_4\tilde p_3 p_3} = \delta_{\tilde p_1 p_2}\sum_{j=0}^{\infty}\sum_{\tilde p_3} P^j_{p_1\tilde p_3\tilde p_3\tilde p_2} \\ &= \frac{\kappa^4}{\lambda^2}\sum_j\sum_{l=0}^{2j}\frac{1}{(2j+1)\big(\gamma(j,l,\alpha,\beta)+\frac{\kappa^4}{\lambda^2}\mu^2\big)}\sum_{\tilde p_3}\sum_k (Y^{j\dagger}_{lk})_{p_1\tilde p_3}(Y^j_{lk})_{\tilde p_3\tilde p_2} \qquad (5.3)\end{aligned}$$

By using the expression of the fuzzy harmonics in the canonical matrix base, (4.40), (4.41), together with the relation

$$\left(\begin{array}{cc|c} j_1 & j_2 & l \\ m_1 & m_2 & k \end{array}\right) = (-1)^{j_1-m_1}\sqrt{\frac{2l+1}{2j_2+1}}\left(\begin{array}{cc|c} j_1 & l & j_2 \\ m_1 & -k & -m_2 \end{array}\right) \qquad (5.4)$$

the sums over $\tilde p_3$ and k can be entirely performed. From the relation

$$\frac{1}{2j+1}(Y^{j\dagger}_{lk})_{p_1\tilde p_3}(Y^j_{lk})_{\tilde p_3\tilde p_2} = (-1)^{-p_1-\tilde p_2}\left(\begin{array}{cc|c} j & j & l \\ \tilde p_3 & -p_1 & k \end{array}\right)\left(\begin{array}{cc|c} j & j & l \\ \tilde p_3 & -\tilde p_2 & k \end{array}\right)$$

we obtain

$$\mathcal{A}_{p_1\tilde p_1;p_2\tilde p_2} = \frac{\kappa^4}{\lambda^2}\delta_{\tilde p_1 p_2}\delta_{p_1\tilde p_2}\sum_{j=0}^{\infty}\sum_{l=0}^{2j}(-1)^{2j}\frac{2l+1}{(2j+1)(\gamma(j,l;\alpha\beta)+\frac{\kappa^4}{\lambda^2}\mu^2)} \qquad (5.5)$$

As it is apparent from (5.5), the amplitude factorizes into a sum over amplitudes at fixed j, each



Figure 3: Planar two-point correlation function

one corresponding to the amplitude on $\mathbb{S}_j$, the fuzzy sphere of radius j.

To compare with known results in the literature on the fuzzy sphere [48, 49] we shall focus on one of these amplitudes at fixed j and further assume $\beta = 0$ in order to reproduce the Laplacian commonly considered. This yields

$$\mathcal{A}^{j\,P}{}_{p_1\tilde{p}_1;p_2\tilde{p}_2} = \frac{\kappa^4}{\lambda^2}\delta_{\tilde{p}_1p_2}\delta_{p_1\tilde{p}_2}\frac{(-1)^{2j}}{2j+1}\sum_{l=0}^{2j}\frac{2l+1}{\alpha l(l+1)+\beta j^2+\frac{\kappa^4}{\lambda^2}\mu^2} \tag{5.6}$$

To move from the interaction base to the propagation base we simply observe that

$$\mathcal{A}^{j\,P} = \sum_{p_i,\tilde{p}_i}\mathcal{A}^{j\,P}{}_{p_1\tilde{p}_1;p_2\tilde{p}_2}\phi^j_{p_1\tilde{p}_1}\phi^j_{p_2\tilde{p}_2} = \sum_{l_ik_i}\tilde{\mathcal{A}}^{j\,P}_{l_1k_1;l_2k_2}\varphi^j_{l_1k_1}\varphi^j_{l_2k_2} \tag{5.7}$$

so that, by means of (4.49) we obtain

$$\tilde{\mathcal{A}}^{j\,P}_{l_1k_1;l_2k_2}(\beta=0) = \frac{\kappa^4}{\lambda^2}\sum_{p_i\tilde{p}_i}\delta_{\tilde{p}_1p_2}\delta_{p_1\tilde{p}_2}\frac{(-1)^{2j}}{2j+1}\sum_{l=0}^{2j}\frac{2l+1}{\alpha l(l+1)+\frac{\kappa^4}{\lambda^2}\mu^2}(Y^j_{l_1k_1})_{p_1\tilde{p}_1}(Y^j_{l_2k_2})_{p_2\tilde{p}_2} \tag{5.8}$$

and using the Kronecker delta symbols, we can perform the sum over $p_i, \tilde{p}_i$. We obtain

$$\tilde{\mathcal{A}}^{j\,P}_{l_1k_1;l_2k_2}(\beta=0) = \frac{\kappa^4}{\lambda^2}\sum_{l}\frac{2l+1}{\alpha l(l+1)+\frac{\kappa^4}{\lambda^2}\mu^2}(-1)^{k_2}\delta_{-k_1k_2}\delta_{l_1l_2} \tag{5.9}$$

which coincides with the result of [48, 49].

As for the full $\mathbb{R}^3_\lambda$ amplitude in the propagating base, we have to restore the term proportional to β and sum over j. We obtain finally

$$\tilde{\mathcal{A}}^{P}_{l_1k_1;l_2k_2} = \frac{\kappa^4}{\lambda^2}\sum_{j\in\mathbb{N}/2}\sum_{l}\frac{2l+1}{\alpha l(l+1)+\beta j^2+\frac{\kappa^4}{\lambda^2}\mu^2}(-1)^{k_2}\delta_{-k_1k_2}\delta_{l_1l_2}. \tag{5.10}$$

We close this subsection by noting that the behavior of the complete amplitude (5.10), restricted to the case $\beta = 0$, is ruled by the behaviour of the series

$$\mathcal{P} = \sum_{n\in\mathbb{N}}\sum_{l=0}^{n}\frac{2l+1}{l(l+1)+\nu^2},\ \nu^2 := \frac{\kappa^4}{\lambda^2}\frac{\mu^2}{\alpha} \tag{5.11}$$

We assume $\nu > 0$. It can be readily observed that (5.11) is divergent. Indeed, let us introduce the following positive function on $[0, +\infty[$, $f(x) = \frac{2x+1}{x(x+1)+\nu^2}$. One can check that it is monotonically decreasing on $[0, +\infty[$ provided $\nu^2 \leq \frac{1}{2}$. Then, from an elementary result of analysis, $\lim_{n\to\infty}\sum_{l=0}^{n}\frac{2l+1}{l(l+1)+\nu^2}$ behaves as $\int_0^\infty dx\ f(x)$. But $\int_0^\infty dx\ f(x) = [\log(x(x+1)+\nu^2)]_0^\infty$ which diverges and so does (5.11). When $\nu^2 \geq \frac{1}{2}$, the above analysis holds true provided one replaces the domain of $f(x)$ by the domain on which $f(x)$ is decreasing and modifies accordingly the lowest value in the summation of the series.

Whenever $\beta \neq 0$, the related complete amplitude (5.10) satisfies $|\tilde{\mathcal{A}}^P_{l_1k_1;l_2k_2}| \leq |\tilde{\mathcal{A}^P}_{l_1k_1;l_2k_2}(\beta = 0)|$. This mixed situation involving both parameters α and β is more complicated and deserves a separate analysis. We postpone to forthcoming publications the detailed analysis of the 1-loop amplitudes for arbitrary values of the parameters together with the perturbative renormalisability of the action (4.1), (4.2) to all orders. The case $\alpha = 0$ yields a finite model and will be considered in the subsection 5.3.

5.2 Non-planar two-point graph

The non-planar contribution to the 2-point correlation function at one loop is represented in Fig. 4. The amplitude is defined as



Figure 4: Nonplanar two-point diagram

$$
\begin{aligned}
\mathcal{A}^{NP}_{p_1\tilde{p}_1;p_2\tilde{p}_2} &= \sum_{j\in\frac{\mathbb{N}}{2}} \sum_{p_3\tilde{p}_3p_4\tilde{p}_4} V^j_{p_1\tilde{p}_1;p_2\tilde{p}_2;p_3\tilde{p}_3;p_4\tilde{p}_4} P^j_{\tilde{p}_4p_4\tilde{p}_2p_2} = \sum_{j=0}^{\infty} P^j_{p_1\tilde{p}_2p_2\tilde{p}_1} \delta_{p_1\tilde{p}_1}\delta_{p_2\tilde{p}_2} \\
&= \frac{\kappa^4}{\lambda^2}\sum_j\sum_{l=0}^{2j} \frac{1}{(2j+1)\big(\gamma(j,l,\alpha,\beta)+\frac{\kappa^4}{\lambda^2}\mu^2\big)} \sum_k (Y^{j\dagger}_{lk})_{p_1\tilde{p}_2}(Y^j_{lk})_{p_2\tilde{p}_1}\delta_{p_1\tilde{p}_1}\delta_{p_2\tilde{p}_2} \\
&= \frac{\kappa^4}{\lambda^2}\sum_j\sum_{l=0}^{2j} \frac{1}{\big(\gamma(j,l,\alpha,\beta)+\frac{\kappa^4}{\lambda^2}\mu^2\big)} \times \\
&\quad \sum_k (-1)^{p_1+\tilde{p}_1} \left(\begin{array}{cc|c} j & j & l \\ \tilde{p}_2 & -p_1 & k \end{array}\right) \left(\begin{array}{cc|c} j & j & l \\ p_2 & -\tilde{p}_1 & k \end{array}\right) \delta_{p_1\tilde{p}_1}\delta_{p_2\tilde{p}_2}
\end{aligned} \tag{5.12}
$$

As before, we can transform this amplitude in the propagation base and compare the case j fixed, $\beta=0$, with [48, 49]. We obtain

$$
\begin{aligned}
&\tilde{\mathcal{A}}^{j\,NP}_{l_1k_1;l_2k_2}(\beta=0) = \mathcal{A}^{j\,NP}_{p_1\tilde{p}_1p_2\tilde{p}_2}(\beta=0)(Y^j_{l_1k_1})_{p_1\tilde{p}_1}(Y^j_{l_2k_2})_{p_2\tilde{p}_2} = \frac{\kappa^4}{\lambda^2}\sum_{l=0}^{2j}\frac{2j+1}{\big(\alpha l(l+1)+\frac{\kappa^4}{\lambda^2}\mu^2\big)} \\
&\sum_{k=-l}^{l}\sum_{p_i,\tilde{p}_i=-j}^{j}(-1)^k \left(\begin{array}{cc|c} j & j & l \\ \tilde{p}_2 & -p_1 & k \end{array}\right) \left(\begin{array}{cc|c} j & j & l \\ p_2 & -\tilde{p}_1 & k \end{array}\right) \left(\begin{array}{cc|c} j & j & l_1 \\ p_1 & -\tilde{p}_1 & k_1 \end{array}\right) \left(\begin{array}{cc|c} j & j & l_2 \\ p_2 & -\tilde{p}_2 & k_2 \end{array}\right) \\
&= \frac{\kappa^4}{\lambda^2}\sum_{l=0}^{2j}\frac{(2j+1)(2l+1)}{\big(\alpha l(l+1)+\frac{\kappa^4}{\lambda^2}\mu^2\big)}(-1)^{l_1+l+2j-k_1}\delta_{l_1l_2}\delta_{k_1,-k_2}\left\{\begin{array}{ccc} j & j & l_1 \\ j & j & l \end{array}\right\}
\end{aligned} \tag{5.13}
$$

where the last term is a Wigner $6j$-symbol. Here we have used the relation between Clebsch-Gordan coefficients and Wigner 3j-symbols

$$
\left(\begin{array}{cc|c} j_1 & j_2 & l \\ p_1 & p_2 & k \end{array}\right) = \sqrt{2l+1}(-1)^{j_1-j_2+k}\begin{pmatrix} j_1 & j_2 & l \\ p_1 & p_2 & -k \end{pmatrix} \tag{5.14}
$$

and the summation formula for the product of four 3j-symbols, as for example in [47]. The result in (5.13) is in agreement with [48, 49].

As for the full $\mathbb{R}^3_\lambda$ amplitude in the propagation base, we restore the term proportional to β and sum the previous expression over j. We obtain

$$\begin{aligned}\tilde{\mathcal{A}}^{NP}_{l_1k_1;l_2k_2} &= \sum_{j\in\frac{\mathbb{N}}{2}} \tilde{\mathcal{A}}^{j\,NP}_{l_1k_1;l_2k_2} \\ &= \frac{\kappa^4}{\lambda^2}\sum_{j\in\frac{\mathbb{N}}{2}}\sum_{l=0}^{2j}\frac{(2j+1)(2l+1)}{\big(\alpha l(l+1)+\beta j^2+\frac{\kappa^4}{\lambda^2}\mu^2\big)}(-1)^{l_1+l+2j-k_1}\delta_{l_1l_2}\delta_{k_1,-k_2}\left\{\begin{array}{ccc} j & j & l_1 \\ j & j & l\end{array}\right\} \end{aligned} \tag{5.15}$$

We now assume $\beta=0$ and study the behavior of the non-planar contribution for small external momenta, i.e $l_1 \ll j$. It is worth pointing out that one expects a behaviour for the full planar and non-planar amplitudes different from what occurs for the fuzzy sphere case [49], which merely coincides with the fixed j contributions. In the present situation, the sum over the sphere of radius j must be taken into account and changes the properties of the amplitudes as we now discuss.

When $l_1=0$, the Wigner $6j$ coefficient in (5.13) can be simplified as

$$\left\{\begin{array}{ccc} j & j & 0 \\ j & j & l\end{array}\right\} = \frac{1}{2j+1}(-1)^{2j+l} \tag{5.16}$$

which, combined with (5.13) yields

$$\tilde{\mathcal{A}}^{NP}_{l_1=0}(\beta=0) = \frac{\kappa^4}{\lambda^2}\sum_{j\in\frac{\mathbb{N}}{2}}\sum_{l=0}^{2j}\frac{(2l+1)}{\big(\alpha l(l+1)+\frac{\kappa^4}{\lambda^2}\mu^2\big)} = \tilde{\mathcal{A}}^{P}_{l_1=0}(\beta=0) \tag{5.17}$$

where $\tilde{\mathcal{A}}^{P}_{l_1=0}(\beta=0)$ can be read off from (5.9). Notice that this relation extends obviously to the case $\beta\neq 0$. From (5.17), one deduces that both planar and non-planar contributions for zero external momentum, $l_1=0$, have the same behavior and are therefore divergent. Notice that $\tilde{\mathcal{A}}^{j,P}_{l_1=0}(\beta=0)=\tilde{\mathcal{A}}^{j,NP}_{l_1=0}(\beta=0)$, for any $j\in\frac{\mathbb{N}}{2}$, therefore recovering a result from [49].

Let us now assume $l_1=1$. By borrowing formulas (3.10) and (3.11) of [49], we obtain for any $j\in\frac{\mathbb{N}}{2}$

$$\tilde{\mathcal{A}}^{j,NP}_{l_1=1,k_1;1,-k_1}(\beta=0) = \frac{\kappa^4}{\lambda^2}(-1)^{k_1}\sum_{l=0}^{2j}\frac{(2l+1)}{\big(\alpha l(l+1)+\frac{\kappa^4}{\lambda^2}\mu^2\big)}\big(1-\frac{l(l+1)}{2j(j+1)}\big) \tag{5.18}$$

and the difference between the fixed j planar and non-planar contributions reads

$$\tilde{\mathcal{A}}^{j,P}_{l_1=1,k_1;1,-k_1}(\beta=0)-\tilde{\mathcal{A}}^{j,NP}_{l_1=1,k_1;1,-k_1}(\beta=0) = \frac{\kappa^4}{\lambda^2}(-1)^{k_1}\sum_{l=0}^{2j}\frac{l(l+1)(2l+1)}{2j(j+1)\big(\alpha l(l+1)+\frac{\kappa^4}{\lambda^2}\mu^2\big)}. \tag{5.19}$$

The RHS of (5.19) is finite and has no singularity which reflects the fact that there is no UV/IR mixing on the fuzzy sphere, in agreement with [49]. However, the difference between the full planar and non-planar amplitudes is divergent. Indeed, from arguments similar to those used at the end of the subsection 5.1, one obtains the following estimate

$$\sum_{n_0}^{2j}\frac{l(l+1)(2l+1)}{2j(j+1)\big(\alpha l(l+1)+\frac{\kappa^4}{\lambda^2}\mu^2\big)} \geq \frac{1}{2j(j+1)}\int_{n_0}^{2j}dx\ \frac{x(x+1)(2x+1)}{\big(\alpha x(x+1)+\frac{\kappa^4}{\lambda^2}\mu^2\big)} \tag{5.20}$$

which holds true from some rank n_0 in the corresponding series. For $j\to\infty$, the RHS of (5.20) behaves as a constant so that

$$\sum_{j\in\frac{\mathbb{N}}{2}}\big(\tilde{\mathcal{A}}^{j,P}_{l_1=1,k_1;1,-k_1}(\beta=0)-\tilde{\mathcal{A}}^{j,NP}_{l_1=1,k_1;1,-k_1}(\beta=0)\big) = +\infty. \tag{5.21}$$

We close this subsection by examining if some UV/IR mixing may occur. This would a priori show up on the 1-loop non-planar amplitude as a singularity in the external momentum l_1, at fixed cut-off on j (recall that the 1-loop planar amplitude has a trivial dependance in l_1). A rigorous analysis would require to make use of asymptotic for the Wigner 6j-symbols. As a first approach to the problem we apply the analysis performed in [49]. On using the following accurate Racah approximation for the Wigner 6j coefficients

$$\left\{ \begin{array}{ccc} j & j & l_1 \\ j & j & l \end{array} \right\} \approx \frac{(-1)^{l_1+l+2j}}{2j} P_{l_1}(1-\frac{l^2}{2j^2}) \tag{5.22}$$

where $P_l(x)$ denotes the Legendre polynomial of order l and is valid provided $l_1 \ll j$ and $j \gg 1$, we write for $j \gg 1$

$$\begin{aligned} \tilde{\mathcal{A}}^{j,NP}_{l_1k_1;l_2k_2}(\beta=0) &- \tilde{\mathcal{A}}^{j,P}_{l_1k_1;l_2k_2}(\beta=0) \\ &\approx (-1)^{k_1}\delta_{k_1,k_2}\delta_{l_1,l_2}\frac{\kappa^4}{\lambda^2}\sum_{l=0}^{2j}\frac{(2l+1)}{\big(\alpha l(l+1)+\frac{\kappa^4}{\lambda^2}\mu^2\big)}(\frac{2j+1}{2j}P_{l_1}(1-\frac{l^2}{2j^2})-1) \\ &\approx (-1)^{k_1}\delta_{k_1,k_2}\delta_{l_1,l_2}\frac{\kappa^4}{\lambda^2}\sum_{l=0}^{2j}\frac{(2l+1)}{\big(\alpha l(l+1)+\frac{\kappa^4}{\lambda^2}\mu^2\big)}(P_{l_1}(1-\frac{l^2}{2j^2})-1). \end{aligned} \tag{5.23}$$

By further assuming that $\mu \ll j$, the sum in (5.23) can be approximated by ($\varepsilon := \frac{1}{j}$.)

$$\begin{aligned} \sum_{l=0}^{2j}\frac{(2l+1)}{\big(\alpha l(l+1)+\frac{\kappa^4}{\lambda^2}\mu^2\big)}(P_{l_1}(1-\frac{l^2}{2j^2})-1) &\approx \frac{1}{\alpha j}\int_0^2 du\ \frac{2u+\varepsilon}{u(u+\varepsilon)+\frac{\mu^2}{j^2}}\big(P_{l_1}(1-\frac{u^2}{2})-1\big) \\ &\approx \frac{1}{2\alpha j}\int_{-1}^{1}\frac{dx}{1-x}(P_{l_1}(x)-1) \\ &= -\frac{1}{\alpha j}h(l_1) \end{aligned} \tag{5.24}$$

where $h(n) := \sum_{k=1}^{n} k^{-1}$, $h(0)=0$ denotes the harmonic number. Eqn. (5.24) simply indicates the absence of UV/IR mixing (up to logarithms) in the scalar field theory restricted to the fuzzy sphere, as found in [49]. Summing over j we conclude

$$\big(\tilde{\mathcal{A}}^{P}_{l_1=1,k_1;1,-k_1}(\beta=0)-\tilde{\mathcal{A}}^{NP}_{l_1=1,k_1;1,-k_1}(\beta=0)\big) \sim h(\Lambda)h(l_1) \tag{5.25}$$

where $h(l_1)$ is finite for small values of l_1. Moreover $h(\Lambda)\sim \log\Lambda$ for large Λ. This indicates a soft behavior in the external momentum and therefore suggests the absence of UV/IR mixing for the present scalar theory on $\mathbb{R}^3_\lambda$.

5.3 A class of finite scalars models at $\alpha = 0$

In this subsection, we set $\alpha = 0$, $\beta \neq 0$ in (4.2). Under this assumption, the propagator (4.55) simplifies into

$$P(\beta)^j_{p_1,\tilde{p}_1;p_2\tilde{p}_2} := (P(0,\beta))^j_{p_1,\tilde{p}_1;p_2\tilde{p}_2} = \frac{\kappa^4}{\lambda^2\beta}\frac{(-1)^{2j}}{j^2+\rho^2}\delta_{\tilde{p}_1p_2}\delta_{\tilde{p}_2p_1},\ \rho^2 := \frac{\kappa^4\mu^2}{\lambda^2\beta},\ \forall j\in\frac{\mathbb{N}}{2} \tag{5.26}$$

which can be obtained by using the properties of the fuzzy harmonics. We will further assume that $\rho \geq 0$ and finite. The complete propagator $P(\beta)_{p_1,\tilde{p}_1;p_2\tilde{p}_2} := \sum_{j\in\frac{\mathbb{N}}{2}} P(\beta)^j_{p_1,\tilde{p}_1;p_2\tilde{p}_2}$ has therefore the following structure

$$P(\beta)_{p_1,\tilde{p}_1;p_2\tilde{p}_2} = \frac{4\kappa^4}{\lambda^2\beta}\sum_{n\in\mathbb{N}}\frac{(-1)^n}{n^2+4\rho^2}\delta_{\tilde{p}_1p_2}\delta_{\tilde{p}_2p_1} \tag{5.27}$$

which simplifies the computation of the amplitude[‖] of any diagram of arbitrary order since each sum over internal indices simply results in an overall factor $2j+1$ while the only remaining sum is over $j \in \frac{\mathbb{N}}{2}$, in view of the expression for the interaction vertex. This will accordingly simplify the power counting analysis to be done in a while.

In order to prepare the ensuing discussion, we now consider the amplitude of the planar diagram for the 2-point function. From (5.5), we find that the corresponding amplitude can be cast into the form

$$\mathcal{A}_{p_1\tilde{p}_1;p_2\tilde{p}_2} = \frac{4\kappa^4}{\lambda^2\beta}\delta_{\tilde{p}_1p_2}\delta_{p_1\tilde{p}_2}\sum_{n=0}^{\infty}(-1)^n\frac{(n+1)}{n^2+4\rho^2} \tag{5.28}$$

It can be shown that $\mathcal{A}_{p_1\tilde{p}_1;p_2\tilde{p}_2}$ is finite. This can be verified for instance by using directly the Dirichlet criterium for convergence of alternate series. Indeed, let us write the sum in (5.28) as

$$\sum_{n=0}^{\infty}(-1)^n\frac{(n+1)}{n^2+4\rho^2} = \sum_{n=0}^{\infty}(-1)^n\frac{n}{n^2+4\rho^2} + \sum_{n=0}^{\infty}(-1)^n\frac{1}{n^2+4\rho^2}.$$

The second series in the RHS obviously satisfies the Dirichlet criterium and therefore is convergent; the first one can be rewritten as

$$\sum_{n=0}^{\infty}(-1)^n\frac{n}{n^2+4\rho^2} = \sum_{n=0}^{[2\rho]}(-1)^n\frac{n}{n^2+4\rho^2} + \sum_{n=[2\rho]+1}^{\infty}(-1)^n\frac{n}{n^2+4\rho^2}$$

where $[z]$ denotes the integer part of z. The first contribution is finite by construction. The second contribution is an alternate series $\sum_n(-1)^n v_n$ with positive decreasing sequence v_n (simply set $v_n = f(n), f(x) = \frac{x}{x^2+4\rho^2}$ is monotonic decreasing on $[2\rho,+\infty[$). It satisfies the Dirichlet test and therefore converges.

For this class of models the analysis of the degree of divergence may be carried out for a generic graph.[**] Let $\mathcal{A}_{\mathcal{D}}$ denote the amplitude for an arbitrary ribbon diagram $\mathcal{D}$ with genus g (g is the genus of the Riemann surface related to the diagram). Recall that a ribbon graph is built from 2 lines (see e.g Fig.1). The relevant topological properties of $\mathcal{D}$ are characterized (see e.g [16]) by a set of integer numbers (V,I,F,B) where V and I denote respectively the number of vertices and internal ribbon lines and the Euler characteristics of the related Riemann surface is $\chi = 2-2g = V-I+F$. F denotes the number of faces which can be determined simply by closing the external legs of $\mathcal{D}$. It counts therefore the number of closed *single* lines. B is the number of boundary components which counts the number of closed lines having external legs. By noting that $F-B$ counts the number of internal summations, we can write (dropping the unessential overall constants)

$$|\mathcal{A}_{\mathcal{D}}| \leq \sum_{j\in\frac{\mathbb{N}}{2}}\left(\frac{1}{j^2+\rho^2}\right)^I(2j+1)^{F-B} \tag{5.29}$$

[‖] Notice that from (5.26), we derive the following upper bound for the complete propagator $|P(\beta)_{p_1,\tilde{p}_1;p_2\tilde{p}_2}| \leq \frac{4!}{\lambda^2\beta}(\zeta(2)+\frac{1}{4\rho^2})$ where $\zeta(z)$ denotes the Riemann ζ-function which however is not strong enough to ensure finitude of the amplitudes

[**] We are indebted with Fabien Vignes-Tourneret for this observation.

The finitude of the RHS of (5.29) depends obviously on the behavior of

$$\sum_{n=0}^{\infty} \frac{(n+1)^{F-B}}{(n^2+4\rho^2)^I} \tag{5.30}$$

which is related to the integral $\int^{\infty} dx\ x^{F-B-2I}$. It is convergent provided

$$\omega(\mathcal{D}) := 2I + B - F = (I + B - 2) + 2g + V > 1 \tag{5.31}$$

which defines the power counting for the noncommutative scalar theory at $\alpha = 0$. This implies that the only potentially divergent diagram of the theory is the one with $g = 0,\ B = I = V = 1$ which corresponds to the planar 2-point function computed just above, which has been found to be finite. This implies that the theory at $\alpha = 0$ is finite.

6 Discussion and conclusion

Let us first summarize the main results of this paper. We have examined the fate of a family of natural scalar NCFT on the noncommutative $\mathbb{R}^3_\lambda$, a deformation of the Euclidean $\mathbb{R}^3$ for which the relevant associative product is not translation-invariant. We have constructed a natural matrix base adapted to $\mathbb{R}^3_\lambda$. It involves the Voros symbols of the operators of the canonical base of $\bigoplus_{j\in\frac{\mathbb{N}}{2}} \mathbb{S}_j$. We have then considered a family of real-valued scalar actions with quartic interaction on $\mathbb{R}^3_\lambda$ whose natural kinetic operator can be written as a linear combination of the square of the angular momentum and a part proportional to the Casimir operator of $\mathfrak{su}(2)$. Working in the natural matrix base, the action can be expressed as an infinite sum of scalar actions defined on the successive fuzzy spheres $\mathbb{S}_j$ that "foliate" the noncommutative space $\mathbb{R}^3_\lambda$, with kinetic operator of Jacobi type. The computation of the propagator in this base, for which the interaction is diagonal, has been done and gives rise to a rather simple expression.

We have computed the planar and non-planar 1-loop contributions to the 2-point correlation function and examined their behavior in two limiting situations. First, when the kinetic operator involves the angular momentum part only, the planar 1-loop contribution is found to be logarithmically divergent. Using the asymptotics of the Wigner $6j$-symbols, we find that the non-planar contribution has a soft behavior in the external momenta (indices). This signals very likely the absence of UV/IR mixing that would destroy the perturbative renormalizability. In the other limit situation with a kinetic operator involving only the radial part, we find that the resulting theory is finite to all orders in perturbation.

Let us discuss the results. When $\alpha \neq 0$ and $\beta = 0$, the soft "low external indices" behavior of the non-planar 2-point function is of logarithmic type since it is related to harmonic numbers. From the dimensional properties of the kinetic operator (see remark below (4.47)), the region with low external indices considered in the subsection 5.2 corresponds naturally to the IR region (i.e low energy excitations from the kinetic operator). As a conclusion, we do not expect that UV/IR mixing spoiling perturbative renormalisability shows up in the corresponding NCFT. As a corollary, we expect that this latter is renormalisable to all orders. The complete proof, including also the general case when α and β are both non zero has been undertaken and will be reported in a forthcoming publication [53]. We notice that the radial term ($\beta \neq 0$) does not play the role of preventing possible occurence of UV/IR mixing, Although this term looks like a Grosse-Wulkenhaar harmonic term, its role is simply to generate a radial dynamics, as can be easily seen from (4.4). The fact that the theory for $\alpha = 0$ and $\beta \neq 0$ is finite to all orders suggests that the addition of the radial term to the angular momentum part in the kinetic operator will likely improve the renormalizability properties of the NCFT. We expect that the extension of the above analysis to the case of noncommutative gauge theories on $\mathbb{R}^3_\lambda$ will give rise to renormalizable noncommutative models of Yang-Mills and Chern-Simons type [31]. We remark finally that the associative product equipping $\mathbb{R}^3_\lambda$ is not invariant by translation. This, combined with the conclusion of this paper about the absence of dangerous UV/IR mixing seems to support the conjecture made in [36], [35] relating translational invariance of the associative product to the possible occurence of troublesome UV/IR mixing. This point must of course be clarified and deserves further investigation.

A Dual Hahn polynomials

Dual Hahn polynomials are in the present framework the counterpart of the Meixner polynomials at the root of the diagonalisation in [16].

Let us consider the expression of the kinetic action in the interaction base (4.19) which we report here for convenience

$$\begin{aligned}(\Delta(\alpha,\beta)+\mu^2\mathbf{1})^j_{m_1\tilde{m}_1;m_2\tilde{m}_2} &= \frac{1}{\lambda^2}\{\delta_{\tilde{m}_1m_2}\delta_{m_1\tilde{m}_2}D^j_{m_2\tilde{m}_2}-\delta_{\tilde{m}_1,m_2+1}\delta_{m_1,\tilde{m}_2+1}B^j_{m_2,\tilde{m}_2}\\ &- \delta_{\tilde{m}_1,m_2-1}\delta_{m_1,\tilde{m}_2-1}H^j_{m_2,\tilde{m}_2}\}\end{aligned} \tag{A.1}$$

with

$$D^j_{m_2\tilde{m}_2} = [(2\alpha+\beta)j^2+2\alpha(j-m_2\tilde{m}_2)]+\lambda^2\mu^2 \tag{A.2}$$

$$B^j_{m_2\tilde{m}_2} = \alpha\sqrt{(j+m_2+1)(j-m_2)(j+\tilde{m}_2+1)(j-\tilde{m}_2)} \tag{A.3}$$

$$H^j_{m_2\tilde{m}_2} = \alpha\sqrt{(j+m_2)(j-m_2+1)(j+\tilde{m}_2)(j-\tilde{m}_2+1)}. \tag{A.4}$$

To find the inverse of (A.1) we go back to the $\mathbb{R}^4$ notation $v^j_{m_i\tilde{m}_i}\to v^j_{p_i\tilde{p}_i}$ with $j+m_i=p_i$, $j+\tilde{m}_i=\tilde{p}_i$, $j-m_i=q_i$, $j-\tilde{m}_i=\tilde{q}_i$ and $i=1,2,p_i\geq 0$ (so that the indices are all positive numbers). The kinetic operator is then represented as

$$\begin{aligned}(\Delta(\alpha,\beta)+\mu^2\mathbf{1})^j_{p_1\tilde{p}_1;p_2\tilde{p}_2} &= \{\delta_{\tilde{p}_1p_2}\delta_{p_1\tilde{p}_2}D^j_{p_2\tilde{p}_2}-\delta_{\tilde{p}_1,p_2+1}\delta_{p_1,\tilde{p}_2+1}B^j_{p_2,\tilde{p}_2}\\ &- \delta_{\tilde{p}_1,p_2-1}\delta_{p_1,\tilde{p}_2-1}H^j_{p_2,\tilde{p}_2}\}\end{aligned} \tag{A.5}$$

with

$$D^j_{p_2\tilde{p}_2} = \beta j^2+2\alpha j(1+p_2+\tilde{p}_2)-2\alpha p_2\tilde{p}_2+\lambda^2\mu^2 \tag{A.6}$$

$$B^j_{p_2\tilde{p}_2} = \alpha\sqrt{(p_2+1)(2j-p_2)(j+\tilde{p}_2+1)(2j-\tilde{p}_2)} \tag{A.7}$$

$$H^j_{p_2\tilde{p}_2} = \alpha\sqrt{p_2(2j-p_2+1)\tilde{p}_2(2j-\tilde{p}_2+1)}. \tag{A.8}$$

We look for orthogonal polynomials which diagonalize the kinetic operator. We pose $\tilde{p}_1-p_1=p_2-\tilde{p}_2=k$ so that

$$(\Delta(\alpha,\beta)+\mu^2\mathbf{1})^j_{p_1\tilde{p}_1;p_2\tilde{p}_2}=(\Delta(\alpha,\beta)+\mu^2\mathbf{1})^j_{p_1,p_1+k;\tilde{p}_2+k,\tilde{p}_2} \tag{A.9}$$

and we look for $U^{(j,k)}_{mi}$ such that

$$(\Delta(\alpha,\beta)+\mu^2\mathbf{1})^j_{p_1,p_1+k;\tilde{p}_2+k,\tilde{p}_2}=\sum_i U^k_{p_1i}(v_i+\mu^2)U^k_{i\tilde{p}_2} \tag{A.10}$$

with $v_i + \mu^2$ the eigenvalues of the kinetic operator and

$$\sum U^{(j,k)}_{mi} U^{(j,k)}_{il} = \delta_{ml}. \tag{A.11}$$

On multiplying on the left by $U^{(j,k)}_{pp_1}$ and summing over p_1 we arrive at

$$U^{(j,k)}_{p\tilde{p}_2}(v_p)(D_{\tilde{p}_2,k+\tilde{p}_2} - v_p - \mu^2) - U^{(j,k)}_{p\tilde{p}_2+1}(v_p) B_{\tilde{p}_2+1,k+\tilde{p}_2+1} - U^{(j,k)}_{p\tilde{p}_2-1}(v_p) H_{\tilde{p}_2-1,k+\tilde{p}_2-1} = 0 \tag{A.12}$$

We redefine

$$U^{(j,k)}_{p\tilde{p}_2}(v_p) = f(p,k,N)\sqrt{\frac{(N-p_2)!p_2!}{\tilde{p}_2!(N-\tilde{p}_2)!}} V^{(N,k)}_{\tilde{p}_2}(v_p) \tag{A.13}$$

with $N = 2j$ and f a normalization factor. We arrive at

$$(D_{\tilde{p}_2 -} v_p - \mu^2) V_{\tilde{p}_2}(v_p) - \alpha(N - \tilde{p}_2)(p_2 + 1) V_{\tilde{p}_2+1}(v_p) - \alpha \tilde{p}_2 (N - p_2 + 1) V_{\tilde{p}_2-1}(v_p) = 0 \tag{A.14}$$

On introducing

$$B_{\tilde{p}_2} = -\alpha(N - \tilde{p}_2)(p_2 + 1) \quad C_{\tilde{p}_2} = -\alpha\, \tilde{p}_2\, (N - p_2 + 1) \tag{A.15}$$

we have

$$B_{\tilde{p}_2} + C_{\tilde{p}_2} = -D_{\tilde{p}_2} + \frac{\beta}{4!}N^2 + \mu^2 \tag{A.16}$$

On redefining

$$\tilde{v}_p = v_p - \frac{\beta}{4!}N^2 - \mu^2 \tag{A.17}$$

we finally obtain

$$\big(-(B_{\tilde{p}_2} + C_{\tilde{p}_2}) - \tilde{v}_p\big) V_{\tilde{p}_2}(v_p) + B_{\tilde{p}_2} V_{\tilde{p}_2+1}(v_p) + C_{\tilde{p}_2} V_{\tilde{p}_2-1}(v_p) = 0 \tag{A.18}$$

This is the equation satisfied by a special class of orthogonal polynomials, the so called *dual Hahn polynomials* [50]

$$V_{\tilde{p}_2}(v_p) \equiv R_{\tilde{p}_2}(\lambda(p); \gamma, \delta, N) \tag{A.19}$$

with the identification $\tilde{v}_p = \lambda(p) = p(p + \gamma + \delta + 1)$, $\gamma = k$, $\delta = -k$.

The dual Hahn polynomials are given in terms of hypergeometric functions as

$$R_n(\lambda(x); \gamma, \delta, N) = {}_3F_2(-n, -x, x + \gamma + \delta + 1; \gamma + 1, -N; 1) \quad 0 \le n \le N \tag{A.20}$$

which are in turn defined in terms of Pochhammer-symbols. When one of the parameters in the first argument of the hypergeometric series is equal to a negative integer $-n$ the series becomes a finite sum, which is our case. We refer to [50] for more details.

We have therefore obtained that the dual Hahn polynomials are the orthogonal polynomials which diagonalize the kinetic part of the action for our scalar model on $\mathbb{R}^3_\lambda$.

Dual Hahn polynomials and fuzzy harmonics are indeed proportional. They are actually well known in nuclear physics and quantum chemistry, were they are also referred to as "discrete spherical harmonics" (see for example [51]).

To clarify the relationship between dual Hahn polynomials and fuzzy harmonics let us reconsider Eq. (4.60) where we map the kinetic action from the interaction to the propagating base, and let us multiply it by $(Y^j_{lk})_{m_1\tilde{m}_1}$. For the sake of clarity we ignore the mass term. Upon summing over $m_1, \tilde{m}_1$ we obtain

$$\begin{aligned} \Delta^j_{m_1\tilde{m}_1 m_2\tilde{m}_2}(Y^j_{lk})_{m_1\tilde{m}_1} &= \frac{1}{2j+1}(-1)^{2j}\delta_{ll_1}\delta_{kk_1}(\Delta^j_{diag})_{l_1k_1l_2k_2}(Y^j_{l_2k_2})_{m_2\tilde{m}_2} \\ &= \frac{1}{\lambda^2}\gamma(j,l;\alpha,\beta)(Y^j_{lk})_{\tilde{m}_2 m_2} \end{aligned} \tag{A.21}$$

which, on inserting the explicit form of $\Delta^j_{m_1\tilde{m}_1m_2\tilde{m}_2}$, and recalling the relation (4.40), can be verified to be the standard recurrence relation for Clebsch-Gordan coefficients

$$\begin{aligned}&[(j-m_2)(j-\tilde{m}_2)(j+m_2+1)(j+\tilde{m}_2+1)]^{\frac{1}{2}}(Y^j_{lk})_{m_2+1,\tilde{m}_2+1}\\ &+[(j+m_2)(j+\tilde{m}_2)(j-m_2+1)(j-\tilde{m}_2+1)]^{\frac{1}{2}}(Y^j_{lk})_{m_2-1,\tilde{m}_2-1}\\ &=[2j(j+1)-l(l+1)-2m_2\tilde{m}_2](Y^j_{lk})_{m_2,\tilde{m}_2},\ \forall j\in\frac{\mathbb{N}}{2},\ -j\leq m_2,\tilde{m}_2\leq j.\end{aligned} \tag{A.22}$$

But, as we have seen in Eq. (A.12), this is also, up to the redefinition (A.13), the recurrence relation for dual Hahn polynomials. To our knowledge this result has first appeared in [52]. The precise relation depends on the normalization chosen for the Hahn polynomials. Up to a normalization function which depends on N, l, k we have

$$R_n(\lambda(l);k,-k,\delta,N)=(-1)^{j-\tilde{m}_2}\sqrt{\frac{(j+\tilde{m}_2)!(j-\tilde{m}_2)!}{(j+m_2)!(j-m_2)!}}(Y^j_{lk})_{m_2\tilde{m}_2} \tag{A.23}$$

with $n=\tilde{p}_2=j+\tilde{m}_2$, $k=m_2-\tilde{m}_2$, $N=2j$.

Acknowledgments: We thank M. Dubois-Violette, H. Grosse, F. Lizzi and F. Vignes-Tourneret for discussions and constructive comments. One of us (P.V.) is grateful to V. Rivasseau for exchanges at various stage of this work. J.-C. W. thanks D. Blaschke and D. Perrot for discussions.

References

[1] A. Connes, *Noncommutative Geometry*, Academic Press Inc., San Diego (1994), available at http://www.alainconnes.org/downloads.html.

[2] For reviews see e.g: A. Connes and M. Marcolli, "*A walk in the noncommutative garden*", (2006), available at http://www.alainconnes.org/downloads.html; G. Landi, "*An introduction to noncommutative spaces and their geometries*", Lectures notes in physics, Springer-Verlag (Berlin, Heidelberg) (1997); J. M. Gracia-Bondía, J. C. Várilly and H. Figueroa, "*Elements of Noncommutative Geometry*", Birkhaüser Advanced Texts, Birkhaüser Boston, Basel, Berlin (2001).

[3] S. Dopplicher, K. Fredenhagen and J. E. Roberts, "Space-time quantization induced by classical gravity "Phys. Lett. B**331** (1994) 39.

[4] E. Witten, "Noncommutative Geometry And String Field Theory", Nucl. Phys. B**268** (1986) 253.

[5] J. Hoppe "Quantum theory of a massless relativistic surface and a two-dimensional bound state problem", Ph.D. thesis, MIT, 1982; published in Soryushiron Kenkyu **80**, 145-202 (1989).

[6] R. L. Stratonovich, "On distributions in representation space". Sov. Phys. JETP **4**, 891 (1957).

[7] J. C. Várilly and J. M. Gracia-Bondía, " The Moyal representation for spin", Ann. Phys. **190**, 107 (1989).

[8] J. Madore, Journ. Math. Phys. **32** (1991) 332; H. Grosse and J. Madore, Phys. Lett. B**283** (1992) 218.

[9] For a review see e.g A. P. Balachandran, S. Kurkcuoglu, S. Vaidya, "Lectures on Fuzzy and Fuzzy SUSY Physics", arXiv:hep-th/0511114 (2005) and references therein.

[10] V. Schomerus, JHEP **9906**, 030 (1999); N. Seiberg and E. Witten, String theory and noncommutative geometry," JHEP 9909 (1999) 032.

[11] L. Susskind, "The Quantum Hall Fluid and Non-Commutative Chern Simons Theory", arXiv:hepth/ 0101029 (2001); S. Hellerman, M. Van Raamsdonk, "Quantum Hall physics equals noncommutative field theory," JHEP **10**, 39-51 (2001); F. Chandelier, Y. Georgelin, T. Masson, J.-C. Wallet, "Quantum Hall conductivity in a Landau type model with a realistic geometry", Ann. Phys. **305** (2003) 60; F. Chandelier, Y. Georgelin, T. Masson, J.-C. Wallet, "Quantum Hall conductivity in a Landau type model with a realistic geometry II", Ann. Phys. **314** (2004) 476.

[12] M. R. Douglas and N. A. Nekrasov, "Noncommutative field theory," Rev. Mod. Phys. **73**, 977 (2001) [hep-th/0106048].

[13] R. J. Szabo, "Quantum field theory on noncommutative spaces," Phys. Rept. **378**, 207 (2003) [hep-th/0109162].

[14] S. Minwalla, M. Van Raamsdonk and N. Seiberg, "Noncommutative perturbative dynamics," JHEP **0002** (2000) 020 [arXiv:hep-th/9912072]; A. Matusis, L. Susskind and N. Toumbas, "The IR/UV connection in the non-commutative gauge theories," JHEP **0012** (2000) 002 [arXiv:hep-th/0002075]; I. Chepelev and R. Roiban, "Renormalization of quantum field theories on noncommutative $\mathbb{R}^d$. I: Scalars," JHEP **0005** (2000) 037 [arXiv:hep-th/9911098].

[15] J.-C. Wallet, "Derivations of the Moyal algebra and Noncommutative gauge theories", SIGMA **5** (2009) 013; E. Cagnache, T. Masson and J-C. Wallet, "Noncommutative Yang-Mills-Higgs actions from derivation based differential calculus“, J. Noncommut. Geom. **5** (2011) 39. A. de Goursac, T. Masson, J.-C. Wallet, "Noncommutative ε-graded connections", J. Noncommut. Geom **6** (2012) 343. For general BRS algebraic algorithms relevant to gauge theories, see e.g J.-C. Wallet, Phys. Lett. B **235** (1990) 71.

[16] H. Grosse and R. Wulkenhaar, "Power-counting theorem for non-local matrix models and renormalisation", Commun. Math. Phys. **254** (2005) 91; H. Grosse and R. Wulkenhaar, "Renormalisation of ϕ^4-theory on noncommutative $\mathbb{R}^2$ in the matrix base", JHEP **0312** (2003) 019; H. Grosse and R. Wulkenhaar, "Renormalisation of ϕ^4-theory on noncommutative $\mathbb{R}^4$ in the matrix base", Commun. Math. Phys. **256** (2005) 305.

[17] M. Buric, J. Madore, "Geometry of the Grosse-Wulkenhaar model", JHEP **1003** (2010) 053. A. Fischer, R. J. Szabo, " Duality covariant quatum field theory on noncommutative Minkowski space", JHEP **0902** (2009) 031. A. de Goursac, A. Tanasa, J.-C. Wallet, "Vacuum configurations for renormalizable noncommutative scalar models", Eur. Phys.J. C**53**(2008)459. A. de Goursac, J.-C. Wallet, "Symmetries of noncommutative scalar field theory", J. Phys. A: Math. Theor. **44** (2011) 055401, [arxiv:0911.2645].

[18] F. Vignes-Tourneret, "Renormalization of the orientable non-commutative Gross-Neveu model," Ann. H. Poincaré **8** (2007) 427. A. Lakhoua, F. Vignes-Tourneret and J. C. Wallet, "One-loop beta functions for the orientable non-commutative Gross-Neveu model," Eur. Phys. J. C **52**, 735 (2007) [arXiv:hep-th/0701170].

[19] C. P. Martin, F. Ruiz Ruiz, "Paramagnetic dominance, the sign of the beta function and UV/IR mixing in non-commutative U(1)", Nucl. Phys. B**597** (2001) 197. M. Attems et al., "Gauge independence of IR singularities in Non-Commutative QFT and interpolating gauges, JHEP **0507** (25) 071. D. N. Baschke, S. Hohenegger and H. Grosse, "Divergences in Non-Commutative gauge theories with Slavnov term“, JHEP **0511** (2005) 041. D. N. Blaschke, H. Grosse, Manfred Schweda, "Non-Commutative U(1) gauge theory on $\mathbb{R} * *4$ with oscillator term and BRST symmetry", Europhys. Lett. **79** (2007) 61002.

[20] A. de Goursac, J. C. Wallet and R. Wulkenhaar, "Noncommutative induced gauge theory," Eur. Phys. J. C **51**, 977 (2007), [arXiv:hep-th/0703075]. H. Grosse and M. Wohlgenannt, "Induced Gauge Theory on a Noncommutative Space," Eur. Phys. J. C **52**, 435 (2007), [arXiv:hep-th/0703169]. A. de Goursac, J.-C. Wallet, R. Wulkenhaar, "On the vacuum states for noncommutative gauge theory", Eur. Phys. J. C **56** (2008) 293, [arxiv:0803.3035].

[21] D. N. Blaschke, A. Rofner, R. I. P Sedmik, M. Wohlgenannt, "On Non-Commutative U*(1) gauge models and renormalisability", J. Phys. A.: Math. Theor. **43** (2010) 425401. D. N. Blaschke, H. Grosse, E. Kronberger, M. Schweda, M. Wohlgenannt,"Loop calculations for the Non-Commutative U(1) gauge field model with oscillator term", Eur. Phys. J. C**67** (2010) 575.

[22] For reviews and references therein , see e.g, J.-C. Wallet, "Noncommutative Induced Gauge Theories on Moyal Spaces," J. Phys. Conf. Ser. **103** (2008) 012007, arXiv:0708.2471 [hep-th]; D. N. Blaschke et

al., “On the problem of renormalisability in Non-Commutative gauge field models - A critical review”, Fortschr. Phys. **58** (2010) 364.

[23] H. Grosse and R. Wulkenhaar, ”The beta-function in duality-covariant noncommutative phi**4 theory”, Eur. Phys. J. C **35**, 277 (2004); M. Disertori, R. Gurau, J. Magnen and V. Rivasseau, ”Vanishing of beta function of non commutative phi(4)**4 theory to all orders”, Phys. Lett. B **649**, 95 (2007).

[24] E. Langmann and R. J. Szabo, ”Duality in scalar field theory on noncommutative phase spaces”, Phys. Lett. B **533** (2002) 168.

[25] H. Grosse and R. Wulkenhaar, ”Self-dual noncommutative φ^4-theory in four dimensions is a non-perturbatively solvable and non-trivial quantum field theory ”, arXiv:1205.0465 (2012).

[26] H. Grosse and R. Wulkenhaar, ”8D-spectral triple on 4D-Moyal space and the vacuum of noncommutative gauge theory,” J.Geom.Phys. 62 (2012) 1583; V. Gayral and R. Wulkenhaar, ”Spectral geometry of the Moyal plane with harmonic propagation”, arxiv:1108.2184 (2011).

[27] J.-C. Wallet, ”Connes distance by examples: Homothetic spectral metric spaces”, Rev. Math. Phys. **24** (2012) 1250027, [arxiv:1112.3285]. See also E. Cagnache, J.-C. Wallet, ”Spectral Distances: Results for Moyal plane and Noncommutative Torus”, SIGMA **6** (2010) 026, [arxiv:0912.4185]; E. Cagnache, F. D’Andrea, P. Martinetti, J.-C. Wallet, ”The spectral distance on the Moyal plane”, J. Geom. Phys. **61** (2011) 1881, [arxiv:0912.0906].

[28] F. D’Andrea, F. Lizzi, J. C. Várilly, “Metric properties of the fuzzy sphere” In press on Lett. Math. Phys. arXiv:1209.0108 [math-ph].

[29] J. M. Gracia-Bondía, F. Lizzi, G. Marmo and P. Vitale, “Infinitely many star products to play with,” *JHEP* **0204**, 026 (2002) [arXiv:hep-th/0112092];

[30] A. B. Hammou, M. Lagraa and M. M. Sheikh-Jabbari, “Coherent state induced star product on R**3(lambda) and the fuzzy sphere,” Phys. Rev. D **66**, 025025 (2002) [arXiv:hep-th/0110291].

[31] P. Vitale, P. Martinetti, J.-C. Wallet, On noncommutative gauge theories on $\mathbb{R}^3_\lambda$, in preparation.

[32] J. M. Gracia-Bondía and J. C. Várilly, “Algebras of distributions suitable for phase space quantum mechanics. I,” J. Math. Phys. **29** (1988) 869; “Algebras of distributions suitable for phase-space quantum mechanics. II. Topologies on the Moyal algebra,” J. Math. Phys. **29** (1988) 880.

[33] A. Voros, “Wentzel-Kramers-Brillouin method in the Bargmann representation”, Phys. Rev. A **40** 6814 (1989).

[34] C. K. Zachos, ”Geometrical evaluation of star products”, J. Math. Phys. 41 (2000) 5129 [arXiv:hep-th/9912238]. G. Alexanian, A. Pinzul and A. Stern, ”Generalized Coherent State Approach to Star Products and Applications to the Fuzzy Sphere,” Nucl. Phys. B**600** (2001) 531 [arXiv:hep-th/0010187].

[35] S. Galluccio, F. Lizzi and P. Vitale, “Twisted Noncommutative Field Theory with the Voros and Moyal Products,” Phys. Rev. D **78** (2008) 085007 [arXiv:0810.2095 [hep-th]].

[36] S. Galluccio, F. Lizzi and P. Vitale, “Translation Invariance, Commutation Relations and Ultraviolet/Infrared Mixing,” JHEP **0909** (2009) 054 [arXiv:0907.3640 [hep-th]].

[37] A. P. Balachandran, A. Ibort, G. Marmo and M. Martone, “Inequivalence of QFT’s on Noncommutative Spacetimes: Moyal versus Voros,” Phys. Rev. D **81**, 085017 (2010) [arXiv:0910.4779 [hep-th]].

A. P. Balachandran and M. Martone, “Twisted Quantum Fields on Moyal and Voros Planes are Inequivalent,” Mod. Phys. Lett. A **24**, 1721 (2009) [arXiv:0902.1247 [hep-th]].

[38] P. Basu, B. Chakraborty and F. G. Scholtz, “A Unifying perspective on the Moyal and Voros products and their physical meanings,” J. Phys. A A **44**, 285204 (2011) [arXiv:1101.2495 [hep-th]].

[39] V. I. Manko, G. Marmo, P. Vitale and F. Zaccaria, “A Generalization of the Jordan-Schwinger map: Classical version and its q deformation,” Int. J. Mod. Phys. A **9**, 5541 (1994) [hep-th/9310053]. For ealier use of Jordan-Schwinger map in various versions of equivariant noncommutative differential calculus, see e.g L.K. Hadjiivanov, R.R. Paunov and I.T. Todorov, J. Math. Phys. **33** (1992) 1379; J.-C. Wallet, J. Phys. A.: Math. Gen. **25** (1992) L1159; Y. Leblanc, J.-C. Wallet, Phys. Lett. B**304** (1993) 89; F. Thuillier, J.-C. Wallet, Phys. Lett. B**323** (1994) 153.

[40] F. Lizzi, P. Vitale and A. Zampini, “The Fuzzy disc,” JHEP **0308** (2003) 057 [hep-th/0306247]. “The

Beat of a fuzzy drum: Fuzzy Bessel functions for the disc," JHEP **0509** (2005) 080 [hep-th/0506008]. "From the fuzzy disc to edge currents in Chern-Simons theory," Mod. Phys. Lett. A **18**, 2381 (2003) [hep-th/0309128].

[41] V. I. Man'ko, G. Marmo and P. Vitale, "Phase space distributions and a duality symmetry for star products" Phys. Lett. A **334**, 1 (2005) [hep-th/0407131]. O. V. Man'ko, V. I. Man'ko, G. Marmo and P. Vitale, "Star products, duality and double Lie algebras," Phys. Lett. A **360**, 522 (2007) [quant-ph/0609041].

[42] A. Tanasa and P. Vitale, "Curing the UV/IR mixing for field theories with translation-invariant $\star$ products," Phys. Rev. D **81**, 065008 (2010) [arXiv:0912.0200 [hep-th]].

[43] F. Lizzi and B. Spisso, "Noncommutative Field Theory: Numerical Analysis with the Fuzzy Disc," Int. J. Mod. Phys. A **27** (2012) 1250137 [arXiv:1207.4998 [hep-th]].

[44] L. Rosa and P. Vitale, "On the $\star$-product quantization and the Duflo map in three dimensions," Mod. Phys. Lett. A **27**, 1250207 (2012) [arXiv:1209.2941 [hep-th]].

[45] G. Marmo, P. Vitale and A. Zampini, "Noncommutative differential calculus for Moyal subalgebras," J. Geom. Phys. **56**, 611 (2006) [hep-th/0411223].

[46] S. R. Das, J. Michelson and A. D. Shapere, "Fuzzy spheres in pp wave matrix string theory," Phys. Rev. D **70** (2004) 026004 [hep-th/0306270].

[47] http://functions.wolfram.com/07.39.23.0014.01

[48] S. Vaidya, "Perturbative dynamics on the fuzzy S**2 and RP**2," Phys. Lett. B **512**, 403 (2001) [hep-th/0102212].

[49] C. -S. Chu, J. Madore and H. Steinacker, "Scaling limits of the fuzzy sphere at one loop," JHEP **0108**, 038 (2001) [hep-th/0106205].

[50] R. Koekoek, P. A. Lesky, R. F. Swarttouw, "Hypergeometric orthogonal polynomials and their q-analogues" , Springer Monographs in Mathematics, Berlin, New York (2010) doi:10.1007/978-3-642-05014-5

[51] V. Aquilanti, S. Cavalli, G. Grossi "Discrete analogs of spherical harmonics and their use in quantum mechanics: the hyperquatization algoritm" Theor. Chim. Acta **79** 283 (1991)

[52] Yu F. Smirnov, S. K. Suslov, A. M. Shirokov, "Clebsch-Gordan coefficients and Racah coefficients for the SU(2) and SU(1,1) groups as the discrete analogues of the Pöschl-Teller potential wavefunctions" J. Phys. A: Math. Gen. **17**, 2157 (1984)

[53] F. Vignes-Tourneret, P. Vitale, J.-C. Wallet, in preparation.